



Thermal pasteurization of ready-to-eat foods and vegetables: Critical factors for process design and effects on quality

Jing Peng, Juming Tang, Diane M. Barrett, Shyam S. Sablani, Nathan Anderson & Joseph R. Powers

To cite this article: Jing Peng, Juming Tang, Diane M. Barrett, Shyam S. Sablani, Nathan Anderson & Joseph R. Powers (2017) Thermal pasteurization of ready-to-eat foods and vegetables: Critical factors for process design and effects on quality, *Critical Reviews in Food Science and Nutrition*, 57:14, 2970-2995, DOI: [10.1080/10408398.2015.1082126](https://doi.org/10.1080/10408398.2015.1082126)

To link to this article: <http://dx.doi.org/10.1080/10408398.2015.1082126>



Accepted author version posted online: 03 Nov 2015.
Published online: 03 Nov 2015.



Submit your article to this journal [↗](#)



Article views: 229



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 2 View citing articles [↗](#)

Thermal pasteurization of ready-to-eat foods and vegetables: Critical factors for process design and effects on quality

Jing Peng^a, Juming Tang^b, Diane M. Barrett^c, Shyam S. Sablani^b, Nathan Anderson^d, and Joseph R. Powers^e

^aCollege of Food Science and Technology, Nanjing Agricultural University, Nanjing, Jiangsu, China; ^bDepartment of Biological Systems Engineering, Washington State University, Pullman, Washington, USA; ^cDepartment of Food Science and Technology, University of California, Davis, California, USA; ^dU.S. Food and Drug Administration, The Center for Food Safety and Applied Nutrition (CFSAN)\Office of Food Safety, Bedford Park, Illinois, USA; ^eSchool of Food Science, Washington State University, Pullman, Washington, USA

ABSTRACT

Increasing consumer desire for high quality ready-to-eat foods makes thermal pasteurization important to both food producers and researchers. To be in compliance with the Food Safety Modernization Act (FSMA), food companies seek regulatory and scientific guidelines to ensure that their products are safe. Clearly understanding the regulations for chilled or frozen foods is of fundamental importance to the design of thermal pasteurization processes for vegetables that meet food safety requirements. This article provides an overview of the current regulations and guidelines for pasteurization in the U.S. and in Europe for control of bacterial pathogens. Poorly understood viral pathogens, in terms of their survival in thermal treatments, are an increasing concern for both food safety regulators and scientists. New data on heat resistance of viruses in different foods are summarized. Food quality attributes are sensitive to thermal degradation. A review of thermal kinetics of inactivation of quality-related enzymes in vegetables and the effects of thermal pasteurization on vegetable quality is presented. The review also discusses shelf-life of thermally pasteurized vegetables.

KEYWORDS

Thermal pasteurization; regulations; pathogens; ready-to-eat; vegetables; FSMA

Introduction

Vegetables are important components for a balanced and healthy diet; they provide essential vitamins, minerals and dietary fiber for our bodies. A range of phytochemicals commonly found in vegetables, such as flavonoids, phenols, and carotenoids, prevent nutritional deficiencies and reduce the risk for various types of cancer, heart disease, diabetes, diverticulosis, stroke, hypertension, birth defects, cataracts, and obesity (Scheerens, 2001; Van Duyn, 1999). However, vegetables are highly perishable and need appropriate preservation technologies to prolong shelf life while maintaining nutritional and sensory qualities.

Emerging alternative technologies are being adopted in addition to traditional thermal processing, thus the term “pasteurization” has been broadened. The requisite scientific parameters for designing a thermal pasteurization process have become more important and need to be clear. Two recently published reviews have provided insights into designing pasteurization processes using 2 particular pathogens in foods as references, namely, nonproteolytic *Clostridium botulinum* spores and *Salmonella* (Silva and Gibbs, 2010, 2012). The reviews by Aamir et al. (2013) and Ling et al. (2015) cover the common and special kinetic models to describe major degradation of food quality attributes during thermal processing. The food industry worldwide has shown strong interest in novel thermal pasteurization processes that can satisfy the increased consumer desire for high quality ready-to-eat meals while

meeting food safety standards. One of the major aims of this review is to highlight the regulations or guidelines associated with food pasteurization by governmental agencies in the U.S. and Europe, and critical factors for designing a pasteurization process accordingly. Current knowledge of food safety regulators and scientists on emerging foodborne pathogens are reviewed, particularly the thermal resistance of several human norovirus surrogates. We will also provide an overview of recent findings on how thermal pasteurization affects vegetable quality, including color, texture, carotenoids, phenolics, antioxidant activity, vitamins and other attributes. In addition, enzyme inactivation and storage conditions which are closely related to the quality of pasteurized vegetables are also discussed here. Technical information provided in this review is related to the thermal processing of fresh and high moisture foods and vegetables.

Development of pasteurization concept

The word “pasteurization” was originally named after the French scientist Louis Pasteur, who invented the process of heating liquids (wine and beer) at a relatively mild temperature (about 55°C) for a short time to prevent spoilage (Silva et al., 2014; Wilbey, 2014). This quick heat method was known as pasteurization and was later applied to milk to kill pathogenic microbes and reduce the number of spoilage organisms. Thus, traditional pasteurization refers to heat

treatment of food (usually below 100°C) to destroy microorganisms of public health significance. Pasteurization processes used in the industry do not kill all microorganisms in foods; they only target pertinent pathogens and reduce spoilage organisms that may grow during storage and distribution (Silva and Gibbs, 2010).

Nowadays pasteurization has been widely accepted as an effective preservation method for killing pathogens in food products, with minimal loss of desired food quality. New technologies that can satisfy the goals of pasteurization have grown rapidly in recent years. The development of these emerging technologies calls for a broadening of the definition of pasteurization. However, there does not appear to be a universally accepted definition for pasteurization. In the United States, the National Advisory Committee on Microbiological Criteria for Foods (NACMCF) has determined the requisite scientific parameters for establishing equivalent alternative methods of pasteurization, and defined pasteurization as “Any process, treatment, or combination thereof that is applied to food to reduce the most resistant microorganism(s) of public health significance to a level that is not likely to present a public health risk under normal conditions of distribution and storage” (NACMCF, 2006). This definition allows application of a broad range of technologies (one or in combination) to different foods for pasteurization, and includes thermal (steam and hot water heating, ohmic heating, microwave heating, infrared processing, etc.) and nonthermal technologies (high pressure processing, ultraviolet radiation, irradiation, pulsed electric field, chemical treatments, ultrasound, filtration, high voltage arc discharge, etc.). In addition to processing method, the following considerations should be included in developing a pasteurization process:

- (1) Determining the most resistant microorganism of public health concern that is likely to survive the selected pasteurization processes;
- (2) Assessing and validating the required level of inactivation of the target microorganism, to make sure it is “not likely to present a public health risk”;
- (3) Evaluating appropriate distribution and storage temperature and shelf life;
- (4) Considering the impact of the food matrix on pathogen survival;
- (5) Defining the specific equipment and operating parameters for the proposed pasteurization process.

Kinetic models and parameters used in design of thermal pasteurization processes

Proper design of industrial thermal pasteurization processes requires knowledge of target microorganism population changes with time at elevated temperatures. It is also desirable to understand thermal impact on important quality attributes. Kinetic models have been widely used to quantify the inactivation of microorganisms and enzymes as well as quality changes in foods caused by thermal processes (Ling et al., 2015; Steinfeld et al., 1998; Van Boekel, 2008). A good starting point for development of kinetic models is

the general rate law:

$$-\frac{dC}{dt} = kC^n, \quad (1)$$

where t is the reaction time, k is the rate constant, n is the order of reaction, and C is the quantity of a chemical compound, quality attribute, enzyme activity or microbial population. Integrating Eq. (1) over time t yields:

$$\text{For } n = 1, C = C_0 e^{-kt}, \quad (2)$$

$$\text{For } n \neq 1, C^{1-n} = C_0^{1-n} + (n-1)kt. \quad (3)$$

C_0 is the initial quantity. The Arrhenius equation is frequently used to describe the temperature effect on the rate constant k of a reaction:

$$k = Ae^{-\frac{E_a}{RT}}, \quad (4)$$

where A is a pre-exponential factor, E_a is activation energy (J/mol), R is the universal gas constant (8.314 J/(mol K)), and T is the absolute temperature (K).

The Arrhenius equation can be alternatively expressed as:

$$k = k_{ref} e^{-\frac{E_a}{R}\left(\frac{1}{T} - \frac{1}{T_{ref}}\right)} \quad (5)$$

where T_{ref} represents the reference temperature (K) and k_{ref} is the reaction rate at the reference temperature.

Two other kinetic parameters, D - and z -values are often used to describe thermal resistance of microorganisms. D -value is the time required at a certain temperature for a one-log reduction (change) in the microorganism population, enzyme activity or quality index; while z -value is the change in temperature required for a 10-fold reduction of the D -value, representing the sensitivity of a reaction to temperature changes.

Thermal kinetic parameters (such as n , k and E_a) in the above equations can be obtained by fitting the models with the experimental data using either one-step or two-step regression methods (Ling et al., 2015). Reaction kinetics in food systems under isothermal conditions (that is at constant temperatures) generally follows a zero, first- or second-order reaction. However, simple reaction models are not always adequate to describe the changes in microorganisms, enzymes or quality in foods. Alternative models such as biphasic first-order, fraction conversion, and Weibull models can provide a better fit (Bozkurt et al., 2014a,b,d; Morales-Blancas et al., 2002).

Kinetic parameters are usually obtained from experiments under isothermal conditions. However, industrial thermal processes are dynamic, in which product temperature changes with time. In order to apply the kinetic data obtained from an isotherm test to predict enzyme inactivation or quality losses during a dynamic thermal process, one can combine the Arrhenius equation with the kinetic model to obtain an integrated equation. This integrated equation can be used to estimate the overall influence of a dynamic thermal process on the activity of an enzyme or quality attributes in foods. Taking the 1st order reaction as an example and substituting alternative Arrhenius

equation Eq. (5) to Eq. (1) ($n = 1$ for the 1st order model) yields:

$$-\frac{dC}{dt} = k_{ref} \left[e^{-\frac{E_a}{R} \left(\frac{1}{T} - \frac{1}{T_{ref}} \right)} \right] C. \quad (6)$$

Integrating both sides of Eq. (6), it becomes:

$$\ln \frac{C(t)}{C_0} = - \int_0^t k_{ref} \left[e^{-\frac{E_a}{R} \left(\frac{1}{T} - \frac{1}{T_{ref}} \right)} \right] dt, \quad (7)$$

where C_0 is the initial quantity for enzyme activity or food quality. The residual enzyme or quality attributes in foods can be calculated by numerical integration of Eq. (7) using the temperature-time history $T(t)$ at a representative location in the food. The representative location should be the least heated spot (cold spot) for enzyme inactivation, the hottest spot (hot spot) for quality degradation. Quality losses can also be calculated for each location and summarized over the entire food volume to assess the overall quality losses.

For food safety, the concept of equivalent time at a reference temperature is used to reflect the cumulative effect of industrial thermal processes. Commercial pasteurization processes are designed to ensure that the accumulated lethal effect at the cold spot in products exceeds a minimum time at a reference temperature specified in regulatory guidance for targeted food pathogens. The equivalent lethal time of a pasteurization process, known as pasteurization value and represented by F , can be calculated by the following equation (FDA, 2011b; Silva and Gibbs, 2010):

$$F = \int_0^t 10^{\frac{T-T_{ref}}{z}} dt, \quad (8)$$

where T ($^{\circ}\text{C}$) is the temperature measured at the cold spot (the area that receives the lowest thermal energy) at time t during a process, T_{ref} is the reference temperature, and z characterizes the sensitivity of thermal resistance of the target bacteria to temperature changes. The reference temperature for pasteurization is usually selected as 70 or 90 $^{\circ}\text{C}$ (ECFF, 2006).

Pathogens of concern for thermal pasteurization

Determining the most resistant microorganism of public health concern for a pasteurization process is of paramount importance. Table 1 lists the primary pathogens of concern and their general growth conditions (ECFF, 2006). Based on Table 1, the optimum refrigerated storage temperature is $\leq 5^{\circ}\text{C}$ (41 $^{\circ}\text{F}$) to eliminate the possibility of growth of most pathogens. However, as much as 10% of product temperatures in consumer refrigerators in the United States are above 7.2 $^{\circ}\text{C}$ (45 $^{\circ}\text{F}$); and temperatures above 10 $^{\circ}\text{C}$ (50 $^{\circ}\text{F}$) for the shelf life of the product would be considered as gross abuse for most refrigerated foods (Audits International, 1999; NACMCF, 2006). Considering the normal distribution and storage temperatures for pasteurized foods with some temperature margin,

pathogens with a minimum growth temperature lower than 7.2 $^{\circ}\text{C}$ (45 $^{\circ}\text{F}$) should be selected as potential pathogens in process design and be included in the hazard analyses. *Listeria monocytogenes*, *Bacillus cereus*, nonproteolytic *Clostridium botulinum*, *Escherichia coli* O157:H7, *Salmonella*, *Staphylococcus aureus*, *Vibrio parahaemolyticus* and *Yersinia enterocolitica* all have minimum growth temperatures of less than 7.2 $^{\circ}\text{C}$ (Table 1). The temperatures to cause one-log reduction of those psychrotrophic microorganisms within 1 min are summarized in Table 2. Spore-forming bacteria *C. botulinum* (nonproteolytic type B) and *B. cereus*, which might cause outbreaks in low-acid under-pasteurized foods, exhibited the highest heat resistance (highest D -values); a temperature less than 85 $^{\circ}\text{C}$ is not adequate to yield even a one-log reduction after several minutes of processing time. But within 1 min at 70 $^{\circ}\text{C}$ was sufficient to achieve a one-log reduction of all the other vegetative bacteria, such as *L. monocytogenes*, *E. coli*, *Salmonella* spp., and *C. burnetii*. Conversely a few seconds at 60 $^{\circ}\text{C}$ is enough to have 90% of *Aeromonas hydrophila* and yeast, which have the lowest heat resistance, inactivated. The thermal resistance of viral pathogens is not included in this table, but is discussed below in Section "Seafood."

Regulations and guidelines for thermal pasteurization in the United States

The Food Safety Modernization Act (FSMA, 2011) requires that preventive controls such as pasteurization be implemented to significantly minimize or prevent the hazards identified in a hazard analysis. Preventive controls will need to be validated. Under the Federal Food Drug and Cosmetic Act (FFDCA, Title 21 section 403(h), 1938), a food shall be deemed to be misbranded as pasteurized unless it is subject to a "safe process," prescribed as pasteurization in a regulation, that is reasonably likely to destroy organisms of public health significance, is at least as effective as the process specified by regulation and is effective throughout the shelf-life of the product when stored under conditions of normal and moderate abuse. Under FFDCA section 403(h), the manufacturer must submit a notification to FDA with data showing effectiveness of the process before labeling a product as "pasteurized."

Table 1. Commonly accepted growth boundaries of pathogenic microorganisms (ECFF 2006).

Microorganism	Min temp ($^{\circ}\text{C}$)	Min pH	Min a_w	Aerobic/ anaerobic
<i>L. monocytogenes</i>	-0.4	4.3	0.92	Facultative
<i>B. cereus</i>	4	4.5	0.93	Facultative
<i>Campylobacter jejuni</i>	32	4.9	0.99	Microaerophilic
<i>C. botulinum</i> (Mesophilic/ proteolytic)	10-12	4.6	0.93	Anaerobic
<i>C. botulinum</i> (Psychrotrophic/ nonproteolytic)	3.3	5.0	0.97 (5% NaCl)	Anaerobic
<i>C. perfringens</i>	12	5.5-5.8	0.935	Anaerobic
<i>E. coli</i>	7-8	4.4	0.95	Facultative
<i>E. coli</i> O157:H7	6.5	4.5	0.95	Facultative
<i>Salmonella</i>	6	4.0	0.94	Facultative
<i>Staphylococcus aureus</i>	5.2	4.5	0.86	Facultative
<i>V. cholera</i>	10	5.0	0.97	Facultative
<i>V. parahaemolyticus</i>	5	4.8	0.94	Facultative
<i>Y. enterocolitica</i>	-1.3	4.2	0.96	Facultative

Table 2. D-values of less than 1 min (0.1–1 min) for psychrotropic microbes under pasteurization temperatures (■ spore-forming pathogens; ■ nonspore forming pathogens; □ nonspore forming spoilage microbes; from Silva and Gibbs (2010) with modification).

Psychrotropic microbes: D-values of < 60 sec at									
55°C	60°C	65°C	70°C	75°C	80°C	85°C	90°C	95°C	>95°C
	<i>Listeria monocytogenes</i>								<i>Bacillus cereus</i>
<i>Escherichia coli</i>							<i>Clostridium botulinum</i> (nonproteolytic type B)		
<i>Salmonella</i> spp.									
	<i>Yersinia enterocolitica</i>								
<i>Aeromonas hydrophila</i>									
	<i>Mycobacterium avium</i>		<i>Coxiella burnetii</i>						
	Lactic acid bacteria								
Yeast									

References in the table include: Ahmed et al. (1995), Bolton et al. (2000), Buduamoako et al., (1992), Cerf & Condron (2006), De-Angelis et al. (2004), Fernández et al. (2001), Franz & vonHoly (1996), Gaze & Brown (1990), Jin et al. (2008), Juneja et al. (1997 & 2007), Juneja & Marmer (1999), Jung & Beuchat (2000), Keswani & Frank (1998), Huang et al. (1992), Miettinen et al. (2005), Murphy et al. (2004a, b), Oteiza et al. (2003), Pearce et al. (2001), Selby et al. (2006), Schuman et al. (1997), Schuman & Sheldon (1997), Thomas et al. (1966), Toora et al. (1992).

At the present, there is no specific guideline available for commercial pasteurization of vegetables in the U.S., because the presence of pertinent bacteria is dependent upon food characteristics, composition (e.g., pH, a_w , salt concentration), and pre- and post-harvest environments; and thermal resistance of bacteria may also vary with different processing technologies. However, there are regulations and standards in place specifically for thermal pasteurization of milk, seafood, egg, and juice products (Table 3) that were promulgated by the responsible governmental authority, either the U.S. Department of Agriculture (USDA) or the U.S. Food and Drug Administration (FDA). It is beneficial to review standards for some of these products.

Milk

Currently, processing of milk is governed by the FDA Pasteurized Milk Ordinance (PMO), and is based on 2 fundamental principles: (1) every particle must be heated to a specified minimum temperature for a specified time and (2) equipment is properly designed and operated. The first federal standard for milk pasteurization was established in 1924, requiring a 61.7°C/30 min process, targeting *Mycobacterium tuberculosis* (Meanwell, 1927). In 1956, *Coxiella burnetii* was recognized as the most resistant bacteria of concern, and the current minimum pasteurization time and temperature combinations (63°C/30 min or 72°C/15 s) were established. Later, Enright (1961) demonstrated a more rigorous pasteurization treatment was needed for 3 milk products: cream, chocolate milk and ice cream mix (Table 3).

Seafood

The pasteurization of seafood is governed by the FDA “Fish and fisheries products hazards and control guidance” (FDA, 2011b). FDA considers a 6D (meaning 6 log reduction) process for *C. botulinum* (type E and nonproteolytic types B and F) to be generally suitable for pasteurized seafood products. In designing a thermal process, a minimum cumulative total lethality of $F_{90^\circ\text{C}}$ (i.e., equivalent accumulated time at 90°C = 10 min) is adequate for pasteurized fish and fishery products (Table 3). For blue crabmeat, the National Blue Crab Industry Pasteurization and Alternative Thermal Processing Standards requires a process of $F_{85^\circ\text{C}} = 31$ min, which exceeds a 12-log

reduction of *C. botulinum* type E spores. Products like Dungeness crabmeat require more rigorous heating ($F_{90^\circ\text{C}} = 57$ min) because it contains certain substances (e.g., lysozyme) that may enable the pathogen to recover more easily after heat damage (FDA, 2011b).

Egg and egg products

Under USDA Food Safety and Inspection Service (FSIS) regulations 9 CFR 590 (CFR, 2012a) the term pasteurize is defined as the “subjecting of each particle of egg products to heat or other treatments to destroy harmful viable microorganisms.” Further this regulation stipulates a “Salmonella negative product” with specific time/temperature requirements given for liquid egg products. FDA regulations 21 CFR 118 (CFR, 2009) stipulate a “...process that achieves at least a 5-log destruction of *Salmonella* for shell eggs, or the processing of egg products in accordance with the Egg Products Inspection Act.” Specific processing conditions for eggs and egg products are listed in Table 3.

Juice

The pasteurization of juice is regulated by the FDA regulation 21 CFR 120, Hazard Analysis and Critical Control Point (HACCP) systems (CFR, 2011). A process with a minimum 5-log reduction of the most resistant microorganism of public health significance identified as the pertinent pathogen under HACCP plan is required. The target microorganisms are dependent on the juice product and process, including *E. coli* O157:H7, *Salmonella*, *Cryptosporidium parvum* or *C. botulinum*. For acidic juices (pH < 4.6), *E. coli* O157:H7, *Salmonella*, and *Cryptosporidium parvum* may occur and cause serious foodborne illness outbreaks; while for low-acid juices such as carrot juice, *C. botulinum* may be present and produce toxins, and therefore becomes the pathogen hazard of concern (CDC, 2006; FDA, 2004).

In summary, the definition of pasteurization in the United States is broad. Regulations and guidelines associated with the pathogens of concern and processing conditions for pasteurization are specific, depending on the particular product, process conditions and packaging systems. There is no “one size fits all” approach to achieving microbiological safety for pasteurization of foods in the United States.

Table 3. Summary of regulations and guidelines for thermal pasteurization of foods (milk, seafood, egg and juice products) in the United States.

Related regulations & guidelines in the U.S.			References
Category I: Milk	The pasteurization of milk is governed by FDA Pasteurized Milk Ordinance. The target microorganism originally was <i>Mycobacterium tuberculosis</i> (1924), now it is <i>Coxiella burnetii</i> (since 1956). A process to eliminate 100, 000 guinea pig infectious doses is needed.		FDA, 2011a
Examples	Target Bacteria	Processing requirements	References
Milk, 1956-present	<i>C. burnetii</i>	63°C (145°F) for 30 min for batch process; 72°C (161°F) for 15 s for HTST	Enright et al., 1957
Cream; Chocolate milk	<i>C. burnetii</i>	66°C (150°F) for 30 min for batch process; 75°C (166°F) for 15 s for HTST	Enright, 1961
Ice cream mix	<i>C. burnetii</i>	69°C(155°F) for 30 min for batch process, 80°C (175°F) for 25 s for HTST	Enright, 1961
Related regulations& guidelines in the U.S.			References
Category II: Seafood	Pasteurization of seafood is governed by FDA (Fish and fisheries products hazards and control guidance). FDA considers a 6D process for target <i>C. botulinum</i> (type E and nonproteolytic types B and F) to be generally suitable for pasteurized seafood products.		FDA, 2011b
Examples	Target bacteria	Processing requirements	References
Fish and fishery products generally (e.g., surimi-based products, soups or sauces)	<i>C. botulinum</i> type E and nonproteolytic types B and F	$F_{90^{\circ}\text{C}} = 10$ min, z value is 7°C for temperatures less than 90°C, 10°C for temperatures above 90°C.	FDA, 2011b
Blue crabmeat	<i>C. botulinum</i> type E and nonproteolytic types B and F	$F_{85^{\circ}\text{C}} = 31$ min, Z value is 9°C.	FDA, 2011b
Dungeness crabmeat	<i>C. botulinum</i> type E and nonproteolytic types B and F	$F_{90^{\circ}\text{C}} = 57$ min, Z value is 8.6°C.	FDA, 2011b
Related regulations& guidelines in the U.S.			References
Category III: Egg products	The current processing of egg products is governed by FSIS regulations 9 CFR 590. A process achieving at least a 5-log reduction of <i>Salmonella</i> is required..		CFR, 2012a
Examples	Products	Processing requirements (Min temp & Min holding time)	References
Liquid eggs	Albumen (w/o use of chemicals)	56.7°C (134°F) for 3.5 min or 55.6 (132°F) for 6.2 min	CFR, 2012b
	Whole egg	60°C (140°F) for 3.5 min	CFR, 2012b
	Whole egg blends (<2% added nonegg ingredients); sugar whole egg (2–12% sugar added); and plain yolk	61.1°C (142°F) for 3.5 min or 60°C (140°F) for 6.2 min	CFR, 2012b
	Fortified whole egg and blends (24–38% egg solids, 2–12% added nonegg ingredients)	62.2°C (144°F) for 3.5 min or 61.1°C (142°F) for 6.2 min	CFR, 2012b
	Salt whole egg (≥2% salt added); sugar yolk (≥2% sugar added); and salt yolk (2–12% salt added)	63.3°C (146°F) for 3.5 min or 62.2°C (144°F) for 6.2 min	CFR, 2012b
Dried egg whites	Spray-dried albumen	54.4°C (130°F) for 7 days	CFR, 2012c
	Pan-dried albumen	51.7°C (125°F) for 5 days	CFR, 2012c
Related regulations& guidelines in the U.S.			References
Category IV: Juice ¹	The pasteurization of juice is governed by FDA (2011, 21 CFR 120.24). A process of 5-log r eduction of most resistant microorganism of public health significance under HACCP plan is required. The target bacteria are dependent on the juice product and process, including <i>E. coli</i> O157: H7, <i>Salmonella</i> , <i>Cryptosporidium parvum</i> or <i>C. botulinum</i> .		CFR, 2011; FDA, 2004
Examples	Target bacteria	References	
Acidic juice (pH ≤ 4.6)	<i>E.coli</i> O157:H7, <i>Salmonella</i> , and <i>Cryptosporidium parvum</i>	FDA, 2004	
Low-acid juices (pH > 4.6)	<i>C. botulinum</i>	FDA, 2004	

¹Juice is defined by FDA as the aqueous liquid expressed or extracted from one or more fruits or vegetables, or concentrates of such liquids or purees.

Regulations and guidelines for thermal pasteurization in Europe

Refrigerated (or chilled) ready-to-eat meals have long been popular in Europe. The European Chilled Food Federation provides guidance for producing chilled foods in Europe (ECFF, 2006). ECFF defines chilled food as “foods that for reasons of safety and/or quality rely on storage at refrigeration

temperatures throughout their entire shelf-life.” According to ECFF recommendations, the common practice for heat-treated chilled food is to aim for a 6 log reduction of either (Table 4):

- (1) *L. monocytogenes* (this treatment will control other vegetative pathogens).
- (2) Cold growing *C. botulinum* (this treatment will not control other spore-forming pathogens such as *B. cereus*).

Table 4. Guidelines for pre-packaged chilled foods (pasteurized foods) in Europe (CSIRO, 2010; ECFF, 2006).

Products	Target bacteria	Processing requirements	Shelf-life
Heat-treated chilled foods ¹	<i>L. monocytogenes</i> <i>Nonproteolytic C. botulinum</i>	6D reduction, common practice of $F_{70^{\circ}\text{C}} = 2.0$ min is considered suitable. 6D reduction, common practice of $F_{90^{\circ}\text{C}} = 10.0$ min is universally accepted	≤ 10 days at 5°C Up to 6 weeks at 5°C

¹Chilled food: foods that for reasons of safety and/or quality rely on storage at refrigeration temperatures throughout their entire shelf life.

L. monocytogenes is the most heat-resistant vegetative pathogen while Type B is the most heat resistant form of nonproteolytic *C. botulinum*. It is generally accepted that a mild pasteurization of low-acid food ($F_{70^{\circ}\text{C}} = 2.0$ min) achieving 6 log reduction of *L. monocytogenes* is suitable for a shelf life of maximally 10 days at 5°C . A severe pasteurization process of $F_{90^{\circ}\text{C}} = 10.0$ min aiming at a 6D process inactivation of nonproteolytic *C. botulinum* allows a product shelf life of up to 6 weeks at 5°C (ECFF, 2006; CSIRO Food and Nutritional Sciences, 2010).

The European Parliament and the Council of the European Union regulation on the hygiene of foodstuffs (EC, 2004) states that for heat treated foods in hermetically sealed containers, any heat treatment process used for an unprocessed product should conform to an internationally recognized standard (e.g., pasteurization) and raise every particle of the product treated to a given temperature for a given period of time.

Apart from the above processing parameters suggested by EFEC for heat-treated chilled foods, commercial production of safe chilled foods should also include the formulation and preservation factors. Variability in the finished product (e.g., pH, a_w) and processing conditions, appropriate hygiene, storage conditions, and usage instructions are important considerations for safety control and in designing the thermal process and assigning shelf-life.

Viral pathogens

Though they are the common cause of foodborne diseases, pathogenic viruses have received far less attention by regulators and in food process design as compared to bacterial pathogens. Foodborne viruses typically are stable outside the host, surviving reasonably well in adverse conditions. Virus transmission can occur by consuming foods that have been contaminated by infected food handlers, during the production process or by the food handling environment. Some viruses associated with food diseases include the human noroviruses (formerly named Norwalk-like viruses), hepatitis A virus (HAV), hepatitis E virus (HEV), rotaviruses, sapoviruses, Aichi virus, astroviruses, adenoviruses, parvoviruses and other small round viruses (Hirneisen et al., 2010; Horm & D'Souza, 2011; Koopmans & Duizer, 2004; Sair et al., 2002). Of these, the noroviruses which cause gastroenteritis, and HAV which causes hepatitis, are currently recognized as the most important 2 foodborne viruses, in terms of the total number of outbreaks and people affected in the Western world. Based on the Centers for Disease Control and Prevention estimates of foodborne illness in the United States, norovirus ranks as the top pathogen contributing to domestically acquired food illnesses, the second resulting in hospitalizations (next to *Salmonella*) and the fourth resulting in death (CDC, 2011). Due to the inability to culture the human norovirus *in vitro*, knowledge about the biology and data on survival of this virus is still limited. No international standard

methods exist for viral analysis of foods, and prevalence studies are limited. The lack of quantitative data on viral load makes the establishment of microbiological criteria for viruses in foods difficult (EFSA, 2011). Current European Food Safety Authority (EFSA) recommendations focus controls on preventive measures to avoid viral contamination rather than trying to inactivate these viruses from food (EFSA, 2011). However, most of the current food hygiene guidelines are optimized for prevention of bacterial infections, and may not be effective against viruses (Koopmans and Duizer, 2004).

Heating appears to be one of the most efficient means for the inactivation of norovirus and hepatitis A & E, despite attempts to use acidification, high pressure processing, chemical disinfection and UV irradiation (Deboosere et al., 2010; Johne et al., 2016). Murine norovirus, feline calicivirus, HAV, canine calicivirus and coliphage MS2 have been used as human norovirus surrogates in various food matrices for survival studies at temperatures relevant to thermal pasteurization. The heat tolerance and survival data of human norovirus surrogates and HAV are summarized in Table 5. These results show that thermal treatment at 60°C for between 1 and 3 min could lead to a one-log reduction of murine norovirus in some tested foods (spinach, raspberry, mussel, milk and water). It also took less than several minutes at 60°C to kill one log of feline calicivirus surrogates based on most reported data (Cannon et al., 2006; Croci et al., 2012). For HAV, most authors found that several minutes of heating at 60°C were sufficient to achieve a one-log decrease in infectivity titer. However, Harlow et al. (2011) reported the *D*-value at 60°C of HAV in mussel to be 109 min; and Gibson and Schwab (2011) reported that the *D*-value at 60°C of HAV in phosphate-buffered saline was 74.6 min. The *z*-values of those viruses fell in the range of 10 – 20°C , as compared to $\sim 10^{\circ}\text{C}$ for bacterial spores (e.g., *C. botulinum*) and $\sim 7^{\circ}\text{C}$ for vegetative bacterial pathogens (e.g., *L. monocytogenes*) (see Table 6).

Thermal inactivation of viruses might be attributed to the changes in the capsid of the virus particle. When the temperature reaches higher than 56 – 60°C , the capsid protein of the virus can be denatured and broken down, which accelerates the inactivation rate of the virus (Croci et al., 1999, 2012; Bozkurt et al., 2013, 2014d). Higher fat and protein content in the heating medium might increase the heat resistance of the virus due to their protective roles (Bidawid et al., 2000; Croci et al., 2012). Other physical and chemical properties of the food matrix (pH, salt and sucrose concentration) may also affect the thermal resistance of the norovirus surrogates (Deboosere et al., 2004; Seo et al., 2012). No ideal human norovirus surrogates have been found so far that would serve all food matrices. For example, canine calicivirus, coliphage MS2 and feline calicivirus are sensitive to low pH while human norovirus is resistant to low pH (pH 2.7 for 3 h at room temperature) (Dolin et al., 1972; Duizer et al., 2004; Cannon et al., 2006; Hewitt



Table 5. Thermal inactivation data associated with foodborne viruses and their surrogates.

	Medium	Temperature range	D or t _b values (min), or log ₁₀ reduction in infectivity titer	Z value (°C)	Ref
Human viruses					
HAV (Hepatitis A virus)	Blue mussel homogenate	50–72°C	D _{50°C} = 54.2, D _{56°C} = 9.32, D _{60°C} = 3.25, D _{65°C} = 2.16, and D _{72°C} = 1.07 min (1st order model); t _b = 37.9, 10.4, 7.73, 6.73 and 1.57 at 50, 56, 60, 65, and 72°C (Weibull model).	15.9° (Weibull models); 13.0 (1 st order models)	Bozkurt et al., 2014a
HAV	Buffered medium	50–72°C	D _{50°C} = 56.2, D _{56°C} = 8.40, D _{60°C} = 2.67, D _{65°C} = 1.73, and D _{72°C} = 0.88 min (1 st order model); t _b = 39.9, 11.1, 4.76, 2.56 and 1.03 min at 50, 56, 60, 65, and 72°C (Weibull model).	14.5 (Weibull models); 12.5 (1 st order models)	Bozkurt et al., 2014b
HAV	Clam	50–72°C	D _{50°C} = 47.4, D _{56°C} = 9.74, D _{60°C} = 6.13, D _{65°C} = 2.20, and D _{72°C} = 1.55 min (1 st order model); t _b = 64.4, 12.6, 5.04, 2.20 and 1.25 min at 50, 56, 60, 65, and 72°C (Weibull model).	13.0 (Weibull models); 14.8 (1 st order models)	Bozkurt et al., 2015b
HAV	Homogenized milk (3.5% fat)	65–85°C	1 log reduction (71°C / 0.18 min); 5 log reduction (80°C / 0.68 min); (Two-phase negative exponential model)	—	Bidawid et al., 2000
HAV	Milk	63 and 72°C	D _{63°C} = 1.1; D _{72°C} ≤ 0.3	—	Hewitt et al., 2009
HAV	Mussel	60–100°C	5 log reduction TCID ₅₀ ml ⁻¹ (100°C / 2min); 4 log reduction TCID ₅₀ ml ⁻¹ (80°C / 10 min)	—	Croci et al., 1999
HAV	Mussel	60–75°C	D _{60°C} = 109; D _{65°C} = 72; D _{70°C} = 17; D _{75°C} = 7	12.0	Harlow et al., 2011
HAV	Mussels au gratin	Cooking water temperature 98°C	<4.6 log reduction TCID ₅₀ ml ⁻¹ (5 min of cooking)	—	Croci et al., 2005
HAV	Mussels hors-d'oeuvre	Cooking water temperature 98°C	<4.6 log reduction TCID ₅₀ ml ⁻¹ (9 min cooking, mussel internal temperature maintained at 71–76°C after the first 3 min)	—	Croci et al., 2005
HAV	Mussels in tomato sauce	Cooking water temperature 98°C	4.6 log reduction TCID ₅₀ ml ⁻¹ (8 min of cooking)	—	Croci et al., 2005
HAV	Phosphate buffered saline	37–70°C	D _{50°C} = 385; D _{60°C} = 74.6; D _{70°C} = 3.84	—	Gibson and Schwab, 2011
HAV	Skim milk (0% fat)	65–85°C	1 log reduction (71°C / 0.16 min); 5 log reduction (80°C / 0.59 min); (Two-phase negative exponential model)	—	Bidawid et al., 2000
HAV	Sodium phosphate buffer	N/A	—	8.40	Harlow et al., 2011
HAV	Spinach	50–72°C	D _{50°C} = 34.4, D _{56°C} = 8.43, D _{60°C} = 4.55, D _{65°C} = 2.30, and D _{72°C} = 0.91 min (1 st order model)	13.9 (1 st order models)	Bozkurt et al., 2015c
HAV	Strawberry mashes	80–90°C	D _{80°C} = 1.22, D _{85°C} = 0.96, D _{90°C} = 0.32 (28° Brix); D _{80°C} = 8.94, D _{85°C} = 4.98, D _{90°C} = 3.00 (52° Brix).	21.4 (28° Brix); 21.1 (52° Brix)	Deboosere et al., 2004
HAV	Synthetic medium (sucrose & Ca)	80–90°C	D _{80°C} = 1.73, D _{85°C} = 0.80, D _{90°C} = 0.22 (28° Brix); D _{80°C} = 12.2, D _{85°C} = 6.28, D _{90°C} = 2.87 (52° Brix).	11.3 (28° Brix); 15.9 (52° Brix)	Deboosere et al., 2004
HAV	Table cream (18% fat)	65–85°C	1 log reduction (71°C / 0.52 min); 5 log reduction (80°C / 1.24 min); (Two-phase negative exponential model)	—	Bidawid et al., 2000
HAV	Turkey deli meat	50–72°C	D _{50°C} = 42.0, D _{56°C} = 20.6, D _{60°C} = 5.9, D _{65°C} = 2.3, and D _{72°C} = 1.0 min (1 st order model); t _b = 25.9, 7.0, 1.7, 0.6 and 0.6 at 50, 56, 60, 65, and 72°C (Weibull model).	13.4 (Weibull models); 12.8 (1 st order models)	Bozkurt et al., 2015a
HAV	Water	63 and 72°C	D _{63°C} = 0.6; D _{72°C} ≤ 0.3	—	Hewitt et al., 2009
Human Norovirus	Spiked mussels	60 and 80°C	less than 1 log RT-PCR units ml ⁻¹ reduction at all temperature-time combinations (up to 15 min)	—	Croci et al., 2012
Human Norovirus	Phosphatase-buffered saline	60 and 80°C	3 log RT-PCR units ml ⁻¹ reduction (80°C / 6 min)	—	Croci et al., 2012

Human virus surrogates															
Canine calicivirus	Modified Eagle's medium	4–100°C	3 D reduction (71.3°C/1 min)												Duizer et al., 2004
Coliphage MS2	Phosphate-buffered saline homogenate	24–85°C	D _{50°C} = 307 (NaCl 0.3%); D _{50°C} = 322 (pH 7) (Weibull models)												Seo et al., 2012
Murine norovirus	Blue mussel homogenate	50–72°C	D _{50°C} = 20.2, D _{56°C} = 6.12, D _{60°C} = 2.64, D _{65°C} = 0.41, and D _{72°C} = 0.18 min (1 st order model); t _D = 19.8, 11.4, 3.01, 0.42 and 0.15 min at 50, 56, 60, 65, and 72°C (Weibull model).												Bozkurt et al., 2014d
Murine norovirus	Buffered medium	50–72°C	D _{50°C} = 36.3, D _{56°C} = 3.74, D _{60°C} = 1.09, D _{65°C} = 0.77, and D _{72°C} = 0.25 min (1 st order model); t _D = 26.8, 2.34, 0.68, 0.39 and 0.09 min at 50, 56, 60, 65, and 72°C (Weibull model).												Bozkurt et al., 2014b
Murine norovirus 1	Dulbecco's Modified Eagle Medium	56 and 73°C	1 log ₁₀ infectivity reduction (56°C/4.21 min) 1 log ₁₀ infectivity reduction (73°C/1.06 min) (Weibull models)												Tuladhar et al., 2012
Murine norovirus	Dulbecco's Modified Eagle Medium	24–85°C	3 log reduction (50°C/3h); 4 log reduction (60°C/10min); 4 log reduction (70°C/2.5 min); D _{50°C} = 54.4 (NaCl 0.3%); D _{50°C} = 123 (pH 7) (Weibull models)												Seo et al., 2012
Murine norovirus	Milk	63 and 72°C	D _{50°C} = 0.7; D _{72°C} = 0.5												Hewitt et al., 2009
Murine norovirus	Modified eagle medium (buffer solution)	50–72°C	D _{50°C} = 34.5, D _{56°C} = 3.65, D _{60°C} = 0.57, D _{65°C} = 0.30, and D _{72°C} = 0.15 min (1 st order model); t _D = 28.3, 3.62, 0.83, 0.37 and 0.11 min at 50, 56, 60, 65, and 72°C (Weibull model).												Bozkurt et al., 2013
Murine norovirus 1	Modified eagle medium	56–72°C	D _{56°C} = 3.47; D _{65°C} = 0.435; D _{72°C} = 0.166												Cannon et al., 2006
Murine norovirus	Phosphate buffered saline	37–60°C	D _{37°C} = 769; D _{50°C} = 106; D _{60°C} = 13.7												Gibson and Schwab, 2011
Murine norovirus 1	Spinach	50–72°C	D _{50°C} = 14.6, D _{56°C} = 3.29, D _{60°C} = 0.98, D _{65°C} = 0.40, and D _{72°C} = 0.16 min (1 st order model); t _D = 15.3, 4.09, 1.11, 0.47 and 0.22 min at 50, 56, 60, 65, and 72°C (Weibull model).												Bozkurt et al., 2014c
Murine norovirus 1	Raspberry puree	65°C/30s or 75°C/15s	1.86 log reduction (65°C/30s); 2.81 log reduction (75°C/15s)												Baert et al., 2008
Murine norovirus 1	1% stool	56 and 73°C	1 log ₁₀ infectivity reduction (56°C/3.20 min)												Tuladhar et al., 2012
Murine norovirus	Turkey deli meat	50–72°C	1 log ₁₀ infectivity reduction (73°C/0.49 min)(Weibull models) D _{50°C} = 21.0, D _{56°C} = 7.3, D _{60°C} = 2.7, D _{65°C} = 0.9, and D _{72°C} = 0.2 min (1 st order model); t _D = 17.8, 6.7, 2.8, 0.9, and 0.3 at 50, 56, 60, 65, and 72°C (Weibull model).												Bozkurt et al., 2015a
Murine norovirus	Water	63 and 72°C	D _{63°C} = 0.9; D _{72°C} ≤ 0.3												Hewitt et al., 2009
Feline calicivirus	Basal Eagle's Medium	56 and 70°C	5–7 log reduction (56°C/60 min)												Doultree et al., 1999
Feline calicivirus	Blue mussel homogenate	50–72°C	3 log reduction (70°C/1 min) D _{50°C} = 5.20, D _{56°C} = 3.33, D _{60°C} = 0.77, D _{65°C} = 0.33, and D _{72°C} = 0.07 min (1 st order model); t _D = 4.03, 3.13, 0.67, 0.31 and 0.08 min at 50, 56, 60, 65, and 72°C (Weibull model).												Bozkurt et al., 2014d
Feline calicivirus	Buffered medium	50–72°C	D _{50°C} = 20.0, D _{56°C} = 6.37, D _{60°C} = 0.94, D _{65°C} = 0.72, and D _{72°C} = 0.21 min (1 st order model); t _D = 13.3, 4.05, 0.40, 0.35 and 0.10 min at 50, 56, 60, 65, and 72°C (Weibull model).												Bozkurt et al., 2014b
Feline calicivirus	Cell culture medium or water	50–75°C	1 log reduction (50°C/15 min); 6 log reduction (70°C/90s)												Buckow et al., 2008
Feline calicivirus	Modified eagle medium (buffer solution)	50–72°C	D _{50°C} = 20.2, D _{56°C} = 6.36, D _{60°C} = 0.56, D _{65°C} = 0.32, and D _{72°C} = 0.11 min (1 st order model); t _D = 13.9, 4.04, 0.37, 0.34 and 0.06 at 50, 56, 60, 65, and 72°C (Weibull model).												Bozkurt et al., 2013

(Continued on next page)

Table 5. (Continued)

	Medium	Temperature range	<i>D</i> or <i>t_D</i> values (min), or \log_{10} reduction in infectivity titer	Z value (°C)	Ref
Feline calicivirus	Modified eagle medium	56–72°C	$D_{56^\circ\text{C}} = 6.72; D_{63^\circ\text{C}} = 0.406; D_{72^\circ\text{C}} = 0.118$	—	Cannon et al., 2006
Feline calicivirus	Modified Eagle's medium	4–100°C	3 D reduction (71.3°C/1 min)	—	Duizer et al., 2004
Feline calicivirus	Phosphate-buffered saline	60 and 80°C	4 log reduction (60°C/3 min) by cell culture assay	—	Croci et al., 2012
Feline calicivirus	Phosphate buffered saline	37–60°C	$D_{37^\circ\text{C}} = 599; D_{50^\circ\text{C}} = 50.6; D_{60^\circ\text{C}} = 14.1$	—	Gibson & Schwab, 2011
Feline calicivirus	Spiked mussels	60 and 80°C	2 log reduction (60°C/15 min) by cell culture assay	—	Croci et al., 2012
Feline calicivirus	Spinach	50–72°C	$D_{50^\circ\text{C}} = 17.4, D_{56^\circ\text{C}} = 5.83, D_{60^\circ\text{C}} = 0.78, D_{65^\circ\text{C}} = 0.27, \text{ and } D_{72^\circ\text{C}} = 0.15$ min (1 st order model); $t_D = 20.7, 6.17, 0.91, 0.35$ and 0.27 min at 50, 56, 60, 65, and 72°C (Weibull model).	10.9 (Weibull models); 9.89 (1st order models)	Bozkurt et al., 2014c
Feline calicivirus	Turkey deli meat	50–72°C	$D_{50^\circ\text{C}} = 9.9, D_{56^\circ\text{C}} = 3.0, D_{60^\circ\text{C}} = 0.8, D_{65^\circ\text{C}} = 0.4, \text{ and } D_{72^\circ\text{C}} = 0.1$ min (1 st order model); $t_D = 11.9, 3.0, 0.8, 0.5$ and 0.1 min at 50, 56, 60, 65, and 72°C (Weibull model).	11.3 (Weibull models); 11.9 (1st order models)	Bozkurt et al., 2015a

^aZ-value for a Weibull model was defined as the change in temperature (°C) required to cause a 90% change in the *t_D*-value of a population (Bozkurt et al., 2013).

^bT(D)₅₀: 50% Tissue Culture Infectious Dose.

Table 6. Temperature-time combinations for “6D process” of some typical bacterial pathogens and viruses under pasteurization.

Target microorganism	Internal product temperature(°C)	Time for 6D process(min)	Lethal rate	Z value(°C)	Reference
<i>Listeria monocytogenes</i>	65	9.3	0.215	7.5	ECFF, 2006
	67	5.0	0.398		
	70	2.0	1.000		
Nonproteolytic <i>Clostridium botulinum</i> type B	90	10.0	1.000	7.0 (T < 90°C)	ECFF, 2006
	93	5.0	2.000	10.0 (T > 90°C)	
	97	2.0	5.010		
Hepatitis A virus	65	13.2	—	14.8	Bozkurt et al., 2015b
	72	9.3	—		
Hepatitis E virus	70	3	—	—	Johne et al., 2016

et al., 2009; Seo et al., 2012). More information on human norovirus (e.g., natural persistence in foods) is needed to refine the regulatory standards and monitoring approaches, and also to develop safe pasteurized foods.

The processing time required to achieve a 6-log reduction of the target pathogen with a typical process time from practical process development view point (in the range of 2–15 min full exposure) for representative pathogens is summarized in Table 6. The most heat-resistant vegetative pathogen, the most heat resistant form of nonproteolytic *C. botulinum* (*C. botulinum* Type B), and the most heat resistant viral pathogen (HAV) from the literature are included in Table 6. Thermal stability of HEV (a popular foodborne virus in Asia, Europe and Africa) is also included. As can be seen, a pasteurization process developed for 6-log reduction of *L. monocytogenes* does not achieve 6-log reductions of HAV or HEV.

Enzyme inactivation in thermal pasteurization

Thermal pasteurization of vegetables aiming to inactivate pathogens may not completely inactivate endogenous enzymes. The presence of residual endogenous enzymes in processed vegetable

products may cause quality loss during storage. Therefore, together with possible growth of spoilage bacteria which in general are more thermally resistant than vegetative pathogenic bacteria, enzyme activity may also considerably shorten the shelf life of the final product. The principal enzyme responsible for a specific quality loss is product-dependent. Major enzymes related to vegetable quality attributes (texture, color and flavor) are listed in Table 7. It can be seen that some enzymes play roles in multiple quality attributes of vegetable products. For example peroxidase (POD) catalyzes the oxidation of phenolics resulting in browning (a color effect), it may also be involved in the oxidative cross-linking of cell wall polymers contributing to the texture changes of vegetables; lipooxygenase (LOX) causes off-flavor through oxidation of polyunsaturated fatty acids, it also catalyzes bleaching or oxidation of carotenoids resulting in the color changes of vegetables. Texture loss through pectin depolymerization can be catalyzed by polygalacturonase (PG), and color loss through browning by POD or polyphenol oxidase (PPO).

Table 8 summarizes published thermal kinetic parameters of several quality-associated enzymes in vegetables. The most frequently studied enzymes are pectin methylesterase (PME), PG, POD, and LOX. Little has been systematically reported on the

Table 7. Major enzymes related to the quality of raw and processed vegetables.

Category	Enzymes	Main effects	Ref
Texture-related enzymes	Pectin methylesterase (PME)	a. Catalyzes the de-esterification of pectin to create binding sites for divalent cations on the polygalacturonic acid backbone of pectin to form cross-links between pectin chains (a firming effect); b. demethoxylated pectin can also be a substrate for PG depolymerization (a softening effect); c. causes cloud loss in juices	Anthon & Barrett, 2002; Terefe et al., 2014; Van Buggenhout, 2009
	Polygalacturonase (PG)	Catalyzes the cleavage of polygalacturonic acid, resulting in pectin depolymerization (softening effect)	
	Peroxidase(POD)	Involved in the oxidative cross-linking of cell wall polysaccharides	
Color-related enzymes	Polyphenol oxidase (PPO)	Acts on phenols in the presence of oxygen, catalyzes browning	Terefe et al., 2014; Zang et al., 2013
	POD	Catalyzes the oxidation of phenolics in the presence of hydrogen peroxide resulting in browning	
	Anthocyanase	Catalyzes the hydrolysis of anthocyanins	
	Chlorophyllase	Catalyzes the degradation of chlorophyll, causes the loss of green color	
Off-flavor-related enzymes	Alliinase	Hydrolyzes the nonprotein amino acid (allicin), involved in the discoloration of processed garlic products	Terefe et al., 2014
	Lipoxygenase (LOX)	Causes the co-oxidation of carotenoids in the presence of free fatty acids, affects the color intensity of foods	
	LOX	Catalyzes the oxidation of polyunsaturated fatty acids, produces volatile off-flavor compounds	
	Hydroperoxidase lyase (HPL)	One of the key enzymes in the “LOX pathway” for producing volatile compounds, the high concentration of which results in off-flavor	
	Cystine lyase	Cleaves cystine producing ammonia, responsible for off flavor and off aroma in broccoli and cauliflower	

PME	Tomato	Citrate buffer, pH 4.5	62–77	1st order	$k_{70^{\circ}\text{C}, \text{R}} = 1.23 \times 10^{-4}$	759 (R)	$D_{70^{\circ}\text{C}} = 5$ (R)	5.01	Guiavar'ch et al., 2003
PME	Tomato	Tris-buffer, pH 7.0	54–81	1st order	$k_{63^{\circ}\text{C}} = 8.30 \times 10^{-2}$; $k_{65^{\circ}\text{C}} = 1.62 \times 10^{-1}$; $k_{67^{\circ}\text{C}} = 3.07 \times 10^{-1}$; $k_{69^{\circ}\text{C}} = 8.80 \times 10^{-1}$	369	$D_{62^{\circ}\text{C}} = 151$; $D_{65^{\circ}\text{C}} = 39.5$ $D_{68^{\circ}\text{C}} = 9.34$; $D_{70^{\circ}\text{C}} = 3.94$	—	Espachs-Barroso et al., 2006
POD	Broccoli (florets)	Phosphate buffer, pH 6.5	70–95	Biphasic 1st order	$k_{70^{\circ}\text{C}, \text{R}} = 4.00 \times 10^{-5}$;	75.0 (L); 58.0 (R)	—	—	Morales-Blancas et al., 2002
POD	Butternut Squash	Butternut squash slices	60–90	Biphasic, 1st order	$k_{65^{\circ}\text{C}, \text{L}} = 2.49 \times 10^1$;	14.0 (L)	—	—	Agüero et al., 2008
POD	Carrot	Carrot juice	70–84	1st order	$k_{65^{\circ}\text{C}, \text{R}} = 7.29 \times 10^{-2}$	15.8 (R)	—	—	Anthón & Barrett, 2002
POD	Carrot	Carrot slices	70–90	1st order	$k_{80^{\circ}\text{C}} = 2.08 \times 10^{-4}$	480	$D_{80.3^{\circ}\text{C}} = 5$	—	Gonçalves et al., 2010
POD	Carrot (cortex)	Phosphate buffer, pH 6.5	70–95	Biphasic 1st order	$k_{80^{\circ}\text{C}} = 4.00 \times 10^{-1}$ $k_{70^{\circ}\text{C}, \text{R}} = 2.10 \times 10^{-6}$;	151.4 95.0 (L); 86.0 (R)	—	—	Morales-Blancas et al., 2002
POD	Carrot (core)	Phosphate buffer, pH 6.5	70–95	Biphasic 1st order	$k_{70^{\circ}\text{C}, \text{L}} = 7.43 \times 10^{-4}$; $k_{95^{\circ}\text{C}, \text{R}} = 7.70 \times 10^{-6}$; $k_{95^{\circ}\text{C}, \text{L}} = 4.80 \times 10^{-3}$;	97.0 (L); 83.0 (R)	—	—	Morales-Blancas et al., 2002
POD	Green asparagus	Green asparagus	70–90	1st order	$k_{70^{\circ}\text{C}, \text{L}} = 1.54 \times 10^{-3}$; $k_{95^{\circ}\text{C}, \text{R}} = 3.30 \times 10^{-6}$; $k_{95^{\circ}\text{C}, \text{L}} = 5.33 \times 10^{-3}$	15.5(bud); 20.7(upper) 23.9 (mid); 22.8 (butt)	—	—	Zheng & Lu, 2011
POD	Green asparagus (tip)	Phosphate buffer, pH 6.5	70–95	Biphasic 1st order	$k_{70^{\circ}\text{C}, \text{R}} = 4.60 \times 10^{-6}$;	67 (L);	—	—	Morales-Blancas et al., 2002
POD	Green asparagus (stem)	Phosphate buffer, pH 6.5	70–95	Biphasic 1st order	$k_{70^{\circ}\text{C}, \text{L}} = 9.58 \times 10^{-4}$; $k_{95^{\circ}\text{C}, \text{R}} = 1.32 \times 10^{-5}$; $k_{95^{\circ}\text{C}, \text{L}} = 4.73 \times 10^{-3}$;	43 (R)	—	—	Morales-Blancas et al., 2002
POD	Potato	Potato homogenate	67–85	1st order	$k_{70^{\circ}\text{C}, \text{L}} = 7.48 \times 10^{-4}$;	53 (R)	—	—	Anthón & Barrett, 2002
POD	Tomato	Tomato slurry	63–67	1st order	$k_{95^{\circ}\text{C}, \text{R}} = 3.84 \times 10^{-5}$; $k_{95^{\circ}\text{C}, \text{L}} = 2.48 \times 10^{-3}$ $k_{80^{\circ}\text{C}} = 5.60 \times 10^{-4}$ $k_{88^{\circ}\text{C}} = 3.32 \times 10^{-2}$; $k_{65^{\circ}\text{C}} = 1.59 \times 10^{-1}$; $k_{67^{\circ}\text{C}} = 6.21 \times 10^{-1}$;	478 149	$D_{83.2^{\circ}\text{C}} = 5$ $D_{63^{\circ}\text{C}} = 69.4$; $D_{65^{\circ}\text{C}} = 14.5$ $D_{67^{\circ}\text{C}} = 3.7$	—	Ercan and Soysal, 2011

(Continued on next page)

Table 8. (Continued)

Enzyme	Source	Heating medium	Temp (°C)	Kinetic model	Kinetic parameters			Ref	
					k (min ⁻¹)	E_a (KJ/mol)	D (min)		
LOX	Broccoli (florets)	Phosphate buffer, pH 6.5	70–95	Biphasic 1st order	$k_{70^\circ\text{C},\text{R}} = 1.15 \times 10^{-6}$; $k_{70^\circ\text{C},\text{L}} = 2.15 \times 10^{-4}$; $k_{95^\circ\text{C},\text{R}} = 4.50 \times 10^{-6}$; $k_{95^\circ\text{C},\text{L}} = 9.10 \times 10^{-4}$; $k_{80^\circ\text{C},\text{R}} = 4.57 \times 10^{-6}$; $k_{80^\circ\text{C},\text{L}} = 1.05 \times 10^{-3}$; $k_{95^\circ\text{C},\text{R}} = 1.15 \times 10^{-5}$; $k_{95^\circ\text{C},\text{L}} = 3.01 \times 10^{-3}$; $k_{80^\circ\text{C},\text{R}} = 1.15 \times 10^{-5}$; $k_{80^\circ\text{C},\text{L}} = 5.83 \times 10^{-4}$; $k_{95^\circ\text{C},\text{R}} = 2.30 \times 10^{-5}$; $k_{95^\circ\text{C},\text{L}} = 1.75 \times 10^{-3}$; $k_{55^\circ\text{C},\text{L}} = 4.21 \times 10^{-2}$; $k_{55^\circ\text{C},\text{R}} = 5.00 \times 10^{-3}$; $k_{65^\circ\text{C},\text{L}} = 1.72$;	—	61 (L); 55 (R)	—	Morales-Blancas et al., 2002
LOX	Green asparagus (tip)	Phosphate buffer, pH 6.5	70–95	Biphasic 1st order		—	76 (L); 65 (R)	—	Morales-Blancas et al., 2002
LOX	Green asparagus (stem)	Phosphate buffer, pH 6.5	70–95	Biphasic 1st order		—	78 (L); 56 (R)	—	Morales-Blancas et al., 2002
LOX	Green bean	Extract solution	55–70	1st order		—	331 (L) 188 (55–63°C, R) 515 (63–70°C, R)	—	Indrawati et al., 1999
LOX	Green pea	Green peas juice	60–72	1st order	$k_{65^\circ\text{C},\text{R}} = 4.80 \times 10^{-2}$; $k_{65^\circ\text{C},\text{L}} = 8.72 \times 10^{-2}$; $k_{70^\circ\text{C}} = 8.02 \times 10^{-1}$	—	541	—	Indrawati et al., 2001
LOX	Green pea	Intact green peas	60–72	1st order	$k_{65^\circ\text{C}} = 1.79 \times 10^{-2}$; $k_{70^\circ\text{C}} = 6.42 \times 10^{-1}$	—	584	—	Indrawati et al., 2001
LOX	Tomato	Tomato dices	80–98	Two-fraction 1st order	$k_{80^\circ\text{C},\text{L}} = 3.00 \times 10^{-2}$; $k_{80^\circ\text{C},\text{R}} = 4.50 \times 10^{-1}$; $k_{98^\circ\text{C},\text{L}} = 2.60 \times 10^{-1}$; $k_{98^\circ\text{C},\text{R}} = 2.21$	$D_{80^\circ\text{C},\text{L}} = 86.9$; $D_{80^\circ\text{C},\text{R}} = 143$ $D_{98^\circ\text{C},\text{L}} = 8.9$; $D_{98^\circ\text{C},\text{R}} = 34.5$	137 (L); 99.7 (R)	18.2 (L) 28.7 (R)	Anese & Sovrano, 2006
LOX	Tomato	Tomato extract	80–98	Two-fraction 1st order	$k_{80^\circ\text{C},\text{L}} = 1.90 \times 10^{-1}$; $k_{80^\circ\text{C},\text{R}} = 1.13$; $k_{90^\circ\text{C},\text{L}} = 7.60 \times 10^{-1}$; $k_{90^\circ\text{C},\text{R}} = 2.00$	$D_{80^\circ\text{C},\text{L}} = 12.2$; $D_{80^\circ\text{C},\text{R}} = 115$ $D_{90^\circ\text{C},\text{L}} = 3.0$; $D_{90^\circ\text{C},\text{R}} = 37.3$	147 (L); 60.8 (R)	16.5 (L) 20.6 (R)	Anese & Sovrano, 2006
LOX	Tomato	Tomato homogenate	50–65	1st order (multiple isoforms)	<i>Isoform 1</i> : $k_{50^\circ\text{C}} = 1.07 \times 10^{-4}$; $k_{65^\circ\text{C}} = 1.67 \times 10^{-2}$; <i>Isoform 2</i> : $k_{50^\circ\text{C}} = 1.50 \times 10^{-5}$; $k_{65^\circ\text{C}} = 4.16 \times 10^{-3}$; <i>Isoform 3</i> : $k_{50^\circ\text{C}} = 1.67 \times 10^{-7}$; $k_{65^\circ\text{C}} = 5.50 \times 10^{-4}$	—	308 (isoform1) 336 (isoform2) 349 (isoform3)	—	Anthon & Barrett, 2003

PG: polygalacturonase; PME: pectin methylesterase; POD: peroxidase; LOX: lipoxygenase; L: heat-labile fraction/form; R: heat-resistant fraction/form.

thermal kinetic parameters of anthocyanase, chlorophyllase, alliinase, or cystine lyase in vegetables. The literature shows that thermal inactivation of PG and PME in vegetables (carrot, tomato and potato) follows a first-order or modified first order (fractional conversion) reaction model. For the inactivation of POD, researchers found that this enzyme system consisted of 2 isozymes with different thermal stability (a heat-labile and a heat-resistant fraction), therefore they applied a biphasic first order model to describe the overall behaviors of this enzyme. This biphasic first order model has been successfully applied to broccoli, butter nut squash, carrot and green asparagus (Morales-Blancas et al., 2002; Agüero et al., 2008). However, other researchers used a monophasic first-order to reflect the kinetic inactivation of POD and also demonstrated a good fit of this model in carrot, green asparagus, potato and tomato (Anthon & Barrett, 2002; Ercan and Soysal, 2011; Gonçalves et al., 2010; Zheng & Lu, 2011).

In terms of the thermal inactivation kinetics of LOX in vegetables, either a biphasic or monophasic first order model was adopted (Anese & Sovrano, 2006; Anthon & Barrett, 2003; Morales-Blancas et al., 2002; Indrawati et al., 2001). Enzymes may have isozymes with different stabilities; and the sequential inactivation pathways may involve intermediate forms with reduced activity, all of which make the enzyme inactivation process complicated. Therefore, a suitable kinetic model should be chosen based on a sufficient understanding of the enzyme (and isozymes) present in a particular food, and all the factors that may affect its activities to accurately predict the enzyme inactivation under thermal processing. The z-values of most reported enzymes listed in Table 8 were below 10°C, except for z-values between 17 and 29°C reported by Anese & Sovrano (2006) for LOX in tomatoes (dices and extract). Based on the publications mentioned above, the required processing time in the range of minutes (<5 min) to achieve 90% inactivation of some quality-related enzymes in vegetables is shown in Table 9. The temperature and time needed to inactivate 90% of the quality-related enzymes depends on the food matrix. Differences in the raw materials and heating medium of an enzyme may result in a wide range of thermal stability. The reported thermal resistance of LOX varied the most, probably due to its different isozymes in addition to the differences in food sources and heating mediums.

Blanching

Blanching with hot water or steam is commonly used as a pre-treatment prior to further processing (canning, freezing, dehydrating etc.) of many vegetables. It is a relatively mild form of heat treatment applied to fresh foods in order to release entrapped air and to inactivate enzymes causing product

quality deterioration (Teixeira, 2014). Typical blanch temperatures and times may be in the range of 190°F (88°C) to 212°F (100°C) for from 1 to 10 min (Barrett, 1994). Low-temperature blanching (generally 50–70°C) prior to high temperature processing has also been studied in a number of vegetables in recent decades (Anthon and Barrett, 2006; Stanley et al., 1995). According to Table 2, several minutes of heating at 70°C may be sufficient for a 6-log reduction in *L. monocytogenes*, which is the most heat-resistant vegetative pathogen. Although the heat treatment applied in blanching has not traditionally been included in microbial inactivation kinetic studies, there are currently research efforts to quantify these and begin to account for them in thermal process applications. Thermal pasteurization of vegetables may or may not involve the blanching step, depending on the specific vegetable products and shelf-life expectations. Additional information about applying blanching prior to vegetable pasteurization, which was obtained from the literature, can be found in Table 10, where both blanching and pasteurization conditions have been specified.

Effect of thermal pasteurization on vegetable quality

Maintaining fresh-like quality is another major goal for vegetable pasteurization, besides microorganism inactivation out of safety concerns. During the thermal pasteurization process, the heat applied to destroy the pathogens also inactivates the endogenous enzymes to limit the quality deterioration of vegetables during processing and storage. The retention of product quality is one factor for determining the shelf-life of pasteurized foods, and also is one main concern for consumer acceptance of processed vegetables. Many papers have reported the impact of thermal pasteurization on the quality of vegetables, including color, texture, carotenoids, phenolics, antioxidant activity, vitamins, and other nutritional attributes. Some of these are summarized in Table 10.

Color

Color plays a vital role in consumer acceptance of a vegetable product, and because we see things first with our eyes before we eat them, it is one of the most important characteristics of vegetables. This visual appeal comes mainly from pigments such as chlorophylls, anthocyanins, and carotenoids including lycopene, which provide health and nutrient benefits. The visual color of vegetables can be numerically expressed by color models. The CIE model is the most commonly used and its 3 color values L^* (lightness), a^* (redness: green to red) and b^*

Table 9. Common processing times (<5 min) to achieve 90% inactivation of some quality-related enzymes in vegetables.

Quality-related enzymes: processing time in minutes to achieve 90% inactivation at									
60°C	65°C	70°C	75°C	80°C	85°C	90°C	95°C	>95°C	
PME ^a		POD ^b			PG ^c				
					LOX ^d				

Adapted from a Anthon & Barrett (2002), Balogh et al. (2004), Espachs-Barroso et al. (2006), Guivarc'h et al. (2003), Ly-Nguyen et al. (2002), Ünal & Bellur (2009); b Anthon & Barrett (2002), Ercan & Soysal (2011); Zheng & Lu (2011); c Anthon & Barrett (2002), Fachin et al. (2003); d Anese & Sovrano (2006), Anthon & Barrett (2002, 2003), Indrawati et al. (1999 & 2001), Morales-Blancas et al. (2002).

Table 10. Effects of thermal pasteurization on the quality of vegetables.

Quality parameters	Commodity/products	Thermal technology	Blanching conditions	Pasteurization conditions	Kinetic parameters	Storage conditions	Enzyme studied	Microbial test	Ref
Antioxidant capacity	Carrot juice	MW heating or traditional heat	—	MW: flow rats 90–287 mL/min at 540–900 W; Traditional heat: 100°C/10 min 90°C/1 min	—	4°C/4 mo	PME	—	Rayman & Baysal, 2011
Antioxidant activity	Gazpacho (a cold vegetable soup)	Tubular heat-exchanger in hot water bath	—	100°C/10 min	—	—	—	—	Elez-Martínez & Martín-Belloso, 2007
Antioxidant activity	Onion by-products (juice, paste and bagasse)	Steam	—	100°C/11–17 min	—	—	—	—	Roldán et al., 2008
Antioxidant capacity	Pumpkin	Thermostatic bath	Boiling water/90 s	85°C/5 min	—	4°C/60 d	—	Total aerobic bacteria, yeast & molds	Zhou et al., 2014
Antioxidant activity	Red sweet pepper	Water bath	—	70°C/10 min	—	—	—	—	—
Antioxidant capacity	HernándezCarrión et al., 2014	Heat exchanger coil in hot water bath	—	90°C/30 or 60 s	$k_{4°C} = 1.37 \times 10^{-2} \text{ d}^{-1}$ (after 90°C/60s heating); $k_{4°C} = 1.22 \times 10^{-2} \text{ d}^{-1}$ (after 90°C/30s heating); 1 st order model.	4°C/91 d	—	—	Odrizola-Serrano et al., 2008
Carotenoids	Carrot juice	MW heating or traditional heat	—	MW: flow rats 90–287 mL/min at 540–900 W; traditional heat: 100°C/10 min	—	4°C/4 mo	—	—	Rayman & Baysal, 2011
Carotenoids	Carrot pieces	Steam	40°C/30 min	$F_{70°C}^{10} = 2\text{min}$; $F_{90°C}^{10} = 10\text{min}$	—	—	—	—	Vervoort et al., 2012
Carotenoids	Red sweet pepper	Water bath	—	70°C/10 min	—	—	—	—	Hernández-Carrión et al., 2014
Carotenoids	Tomato juice	Water bath	—	90°C/30 or 60 s	—	4±1°C/56 d	—	—	Odrizola-Serrano et al., 2009
β-carotene	Carrot pieces	Steam	60°C water/40 min	Actual $F_{70°C} = 1.85$ min or $F_{90°C} = 9.67$ min	—	—	—	—	Lemmens et al., 2013
β-carotene	Carrot puree	Steel tubes in water bath	—	$F_{90°C} = 10\text{min}$	—	—	—	—	Knockaert et al., 2012
Color	Asparagus	Water bath	—	70–98°C	$E_a = 13.1 \text{ kcal/mol}$, $k_{84°C} = 6.60 \times 10^{-3} \text{ min}^{-1}$, 1 st order model	—	—	—	Lau et al., 2000
Color	Broccoli (florets & stems)	Continuous microwave	95°C water/30 s	3.5 kW/4 min, surface temp. of vegetable packs 75–80°C.	—	30°C/60 d	—	Visual signs of spoilage, bacterial and yeast growth	Koskinemi et al., 2013
Color	Carrot pieces	Steam	40°C/30 min	$F_{70°C}^{10} = 2$ min; $F_{90°C}^{10} = 10$ min	—	—	—	—	Vervoort et al., 2012

Color	Cucumber juice	Heat exchanger	85°C/ 15 s	—	4 ± 2°C/ 50d	LOX	Total aerobic bacteria	Zhao et al., 2013
Color	Pickled garlic	Hot water bath	90°C water/ 15 min	—	27 ± 2°C/ 4mo	—	Enterobacteriaceae, lactic acid bacteria & yeast	Rejano et al., 1997
Color	Pumpkin	Thermostatic bath	85°C/ 5 min	—	4°C/ 60 d	—	Total aerobic bacteria, yeast & molds	Zhou et al., 2014
Color	Red bell pepper	Continuous microwave	3.5 kW/ 4 min, surface temp. of vegetable packs 75–80°C.	—	30°C/ 60 d	—	Visual signs of spoilage, bacterial and yeast growth	Koskinieni et al., 2013
Color	Soybean sprouts	Heat (no information on the heating media)	F _{90°C} = 10 min; F _{70°C} = 2 min	—	3°C/ 36 d; 10°C/ 24 d	—	Aerobic, anaerobic & psychrophilic bacterial counts	Koo et al., 2008
Color	Spinach	Oil bath	65–90°C	—	—	—	—	Aamir et al., 2014
Color	Sweatpotato	Continuous microwave	3.5 kW/ 4 min, surface temp. of vegetable packs 75–80°C.	—	30°C/ 60 d	—	Visual signs of spoilage, bacterial and yeast growth	Koskinieni et al., 2013
Color	Tomato juice	Water bath	90°C/ 30 or 60 s	—	4 ± 1°C/ 56 d	—	—	Odrozola-Serrano et al., 2009
Color	Vegetable juice	Oil batch	80–100°C/ 0–60 s	—	—	—	—	Loong and Goh, 2004
Lycopene	Tomato juice	Heat exchanger coil in hot water bath	90°C/ 30–60 s	—	4°C/ 91 d	—	—	Odrozola-Serrano et al., 2008

(Continued on next page)

a: zero-order; b: zero-order for summer and 1st order for the rest seasons
 $k_{90°C} = 1.33 \times 10^{-2}$;
 $k_{100°C} = 1.75 \times 10^{-2}$ (L value, s⁻¹);
 $k_{80°C} = 1.14 \times 10^{-2}$;
 $k_{90°C} = 1.81 \times 10^{-2}$;
 $k_{100°C} = 2.80 \times 10^{-2}$ (a value, s⁻¹); 1st order model.
 $k_{4°C} = 1.77 \times 10^{-2}$ d⁻¹ (after 90°C/ 60s heating);
 $k_{4°C} = 2.27 \times 10^{-2}$ d⁻¹ (after 90°C/ 30s heating); 1st order model.



Table 10. (Continued)

Quality parameters	Commodity/products	Thermal technology	Blanching conditions	Pasteurization conditions	Kinetic parameters	Storage conditions	Enzyme studied	Microbial test	Ref
Phenolics	Carrot juice	MW heating or traditional heat	—	MW: flow rats 90–287 mL/min at 540–900 W; Traditional heat: 100°C/10 min	—	4°C/4 mo	—	—	Rayman & Baysal, 2011
Phenolics	Onion by-products	Steam	—	100°C/11–17 min	—	—	—	—	Roldán et al., 2008
Phenolics	Tomato juice	Heat exchanger coil in hot water bath	—	90°C/30 or 60 s	—	4°C/91 d	—	—	Odrozola-Serrano et al., 2008
Phenolics	Tomato juice	Water bath	—	90°C/30 or 60 s	—	4±1°C/56 d	—	—	Odrozola-Serrano et al., 2009
Phenols (total)	Pumpkin	Thermostatic bath	Boiling water/90 s	85°C/5 min	—	4°C/60 d	—	Total aerobic bacteria, yeast & molds	Zhou et al., 2014
Quercetin	Onion by-products	Steam	—	100°C/11–17 min	—	—	—	—	Roldán et al., 2008
Quercetin	Yellow "banana" pepper	Heat (no information on the heating media)	—	74°C/10 min	—	23°C/124 d	—	—	Lee & Howard, 1999
Quercetin	Tomato juice	Water bath	—	90°C/30 or 60 s	—	4±1°C/56 d	—	—	Odrozola-Serrano et al., 2009
Texture	Asparagus	Water bath	—	70–98°C	$E_a = 24 \text{ kcal/mol}$; $k_{84^\circ\text{C}} = 1.60 \times 10^{-2} \text{ min}^{-1} \text{ (butt)}$; $k_{84^\circ\text{C}} = 2.70 \times 10^{-2} \text{ min}^{-1} \text{ (bud)}$; 1 st -order model	—	—	—	Lau et al., 2000
Texture	Asparagus	Microwave or hot water	—	88°C/10 s	—	—	POD (final products)	—	Lau & Tang, 2002
Texture	Broccoli	Continuous microwave	95°C water/30 s	3.5 kW/4 min, surface temp. of vegetable packs 75–80°C.	—	30°C/60 d	—	Visual signs of spoilage, bacterial and yeast growth	Koskinemi et al., 2013
Texture	Carrot dices	Oil bath	—	80–110°C	$E_a = 140 \text{ kJ/mol}$ in distilled water; 2 nd order model	—	—	—	Peng et al., 2014
Texture (Total pectin contents)	Carrot juice	MW heating or traditional heat	—	MW: flow rats 90–287 mL/min at 540–900 W; traditional heat: 100°C/10 min	—	4°C/4 mo	PME	—	Rayman & Baysal, 2011
Texture	Carrot pieces	Steam	40°C/30 min	$F_{70^\circ\text{C}}^{10} = 2 \text{ min}$ or $F_{90^\circ\text{C}}^{10} = 10 \text{ min}$	—	—	PME & POD (final product)	—	Vervoort et al., 2012
Texture	Carrot pieces	Steam	60°C water/40 min	Actual $F_{70^\circ\text{C}} = 1.85 \text{ min}$ or $F_{90^\circ\text{C}} = 9.67 \text{ min}$	—	—	—	—	Lemmens et al., 2013
Texture	Jalapeño pepper rings	Water (preheating), Steam (heating)	50°C/0–60 min	75°C/5 min	—	23°C/5 mo	—	—	Howard et al., 1997
Texture	Pickled garlic	Hot water bath	90°C water/15 min	90°C/5.5 min	—	27±2°C/4 mo	—	Enterobacteriaceae, lactic acid bacteria & yeast	Rejano et al., 1997

Texture	Pumpkin	Thermostatic bath	Boiling water/ 90 s	85°C/ 5 min	—	4°C/ 60 d	—	Total aerobic bacteria, yeast & molds	Zhou et al., 2014
Texture	Red bell pepper	Continuous microwave	95°C water/ 30 s	3.5 kW/ 4 min, surface temp. of vegetable packs 75–80°C	—	30°C/ 60 d	—	Visual signs of spoilage, bacterial and yeast growth	Koskinemi et al., 2013
Texture	Red sweet pepper	Water bath	—	70°C/ 10 min	—	—	—	—	Hernández- Carrión et al., 2014 Koo
Texture	Soybean sprouts	Heat (no information on the heating media)	100°C steam/ 8 min	$F_{90°C} = 10$ min or $F_{70°C} = 2$ min	—	3°C/ 36 d; 10°C/ 24 d	—	Aerobic, anaerobic & psychrophilic bacterial counts	Koo et al., 2008
Texture	Sweet potato	Continuous microwave	95°C water/ 30 s	3.5 kW/4 min, surface temp. of vegetable packs 75–80°C	—	30°C/ 60 d	—	Visual signs of spoilage, bacterial and yeast growth	Koskinemi et al., 2013
Texture	Yellow "banana" pepper	Heat (no information on the heating media)	—	74°C/ 10 min	—	23°C/ 124 d	—	—	Lee & Howard, 1999
Vitamin C	Gazpacho (a cold vegetable soup)	Tubular heat- exchanger in hot water bath	—	90°C/ 1 min	—	—	—	—	Elez-Martínez & Martín- Belloso, 2007
Vitamin C	Orange-carrot juice	Plate heat exchanger	—	98°C/ 21 s	$k_{10°C} = -9.63 \times 10^{-2} d^{-1}$; $k_{2°C} = -4.66 \times 10^{-2} d^{-1}$; 1st order model.	2°C/ 70 d; 10°C/ 59 d	—	—	Torregrosa et al., 2006
Vitamin C	Pumpkin	Thermostatic bath	Boiling water/ 90 s	85°C/ 5 min	—	4°C/ 60 d	—	Total aerobic bacteria, yeast & molds	Zhou et al., 2014
Vitamin C	Soybean sprouts	Heat (no information on the heating media)	100°C steam/ 8 min	$F_{90°C} = 10$ min; $F_{70°C} = 2$ min	—	3°C/ 36 d; 10°C/ 24 d	—	Aerobic, anaerobic & psychrophilic bacterial counts	Koo et al., 2008
Vitamin C	Yellow "banana" pepper	Heat (no information on the heating media)	—	74°C/ 10 min	—	23°C/ 124 d	—	—	Lee & Howard, 1999
Vitamin C	Tomato juice	Heat exchanger coil in hot water bath	—	90°C/ 30 or 60 s	$k_{4°C} = 3.04 \times 10^{-2} d^{-1}$ (after 90°C/ 60s heating); $k_{4°C} = 3.75 \times 10^{-2} d^{-1}$ (after 90°C/ 30s heating); 1st order model.	4°C/ 91 d	—	—	Odirozola- Serrano et al., 2008
Vitamin D	Vegetable juice	Plate heat exchanger	—	90°C/ 15 s	—	—	—	—	Barba et al., 2012
Vitamin E	Vegetable juice	Plate heat exchanger	—	90°C/ 15 s	—	—	—	—	Barba et al., 2012

(yellowness: blue to yellow) can be used individually or in combination in the form of hue ($h_{ab} = \arctan \frac{b^*}{a^*}$), chroma ($C_{ab}^* = \sqrt{a^{*2} + b^{*2}}$), or total color difference value ($\Delta E = \sqrt{\Delta L^{*2} + \Delta a^{*2} + \Delta b^{*2}}$).

Publications related to color changes of vegetables by thermal pasteurization are listed in Table 10. Most authors found that color degradation in vegetables by thermal pasteurization depends mainly on the heat intensity, duration, media, compounds responsible for color, and storage time. For example, Koskiniemi et al. (2013) pasteurized 3 vegetables (broccoli, red bell pepper and sweet potato), and found that the green color of broccoli florets changed the most while the sweet potato color was stable over the course of processing. Kinetic models have been developed to predict the changes of vegetable colors during the pasteurization process. The first order kinetic model was successfully applied to describe the color degradation in asparagus at temperatures ranging from 70 to 98°C, and in a mixed vegetable juice (butterhead lettuce, celery, parsley, apple concentrate and kalamansi lime) between 80 and 100°C (Lau et al., 2000; Loong and Goh, 2004). Both studies referred to a decrease in green color, which closely correlated with reduction of chlorophyll content in the products. Aamir et al. (2014) indicated that the color changes in spinach leaves followed a zero- or first-order kinetic model under pasteurization conditions, depending on the color parameters measured and the harvest season of the spinach. Among previous studies, 6 included the study of color changes during the storage period. The discoloration of pressurized vegetables and vegetable products can occur due to enzymatic or nonenzymatic browning. The desired color values of most pasteurized vegetables decreased during the storage period, which varied from 36 to 120 days in these studies (Koo et al., 2008; Koskiniemi et al., 2013; Rejano et al., 1997; Zhao et al., 2013; Zhou et al., 2014). Only one study by Koskiniemi et al. (2013) reported that the color of broccoli florets did not change during an extended storage or even during thermal processing. The reason was that addition of acid during the equilibration process had a profound effect on the broccoli color, converting chlorophyll (green) into pheophytin (olive green); processing and storage did not produce further changes in the measured color component. The conversion of chlorophyll to pheophytin and pheophorbide can be accelerated when heating green vegetables under lower pH (acidic) conditions, which results in the discoloration of green vegetables during processing (Andrés-Bello et al., 2013).

Carotenoids

Carotenoids are one of the predominant organic pigments present in vegetables, and include α - and β -carotenes (yellow/orange), lycopene (red/orange), xanthophyll (yellow), lutein, and zeaxanthin (green/yellow). They are also one of the important bioactive compounds in carrots, and can act as antioxidants to reduce the risk of developing degenerative diseases. For example, α - and β -carotenes, vitamin A precursors responsible for the orange color in vegetables such as carrot and sweet potato, are also important for vision. Lycopene is considered a potential antioxidant and cancer-preventing agent, responsible for the red color in tomatoes. Lycopene, α - and β -carotenes

may undergo isomerization, oxidation and other chemical changes during thermal processing and storage due to their highly unsaturated structure (Rodríguez-Amaya and Kimura, 2004; Shi et al., 2003).

The effect of thermal pasteurization on the carotenoids in vegetables depends on the heat intensity and the properties of the products. Total carotenoids found in vegetables are relatively stable to mild pasteurization. Vervoort et al. (2012) heated carrot pieces under conditions from mild pasteurization ($F_{70^\circ\text{C}} = 2$ min) to severe pasteurization ($F_{90^\circ\text{C}} = 10$ min) and found no considerable differences occurred in total carotenoid content after processing, or their individual α - or β -carotene concentration. The authors attributed the stability of carotenes to the protective food matrix, which preserves them from degradation during pasteurization; the applied pasteurization conditions were not severe enough to cause a notable isomerization and/or oxidation of the carotenoids in carrots. Similar results were obtained by Lemmens et al. (2013), for β -carotene content in carrots. However, Odriozola-Serrano et al. (2009) observed an increase of total carotenoid content, lycopene and β -carotene after pasteurization of tomato juice at 90°C for 30 or 60 s. One explanation given was that the homogenization and heat treatment conditions disrupted cell membranes and protein-carotenoid complexes, increasing the extractability of the carotenoids. The carotenoid content of vegetables was reported as decreasing after pasteurization in some studies. Rayman and Baysal (2011) reported a decrease in total carotenoid content of carrot juice after pasteurization at 100°C for 10 min, and Hernández-Carrión et al. (2014) observed a significant reduction of carotenoids in red sweet pepper after pasteurization at 70°C for 10 min. However, the latter study reported the carotenoid content in red sweet pepper on a wet weight basis, which reduces the reliability of their results.

Carotenoids in processed vegetables in storage may also undergo oxidation and isomerization of the all-trans form to the cis forms, causing changes in their levels in the final products. A decreasing trend of total carotenoid content and lycopene content in pasteurized tomato juice was observed by Odriozola-Serrano et al. (2008, 2009) during 3 months storage at 4°C. The authors explained that the decrease in lycopene content throughout storage may be due to the oxygen available in the headspace of the container. In contrast, Rayman and Baysal (2011) reported an increase in the carotenoid content in carrot juice after 3 months of storage at 4°C, and attributed it to possible isomerization of β -carotene.

Thermal pasteurization may also influence the bioaccessibility of carotenoids of vegetables. The bioaccessibility refers to the fraction of a nutrient that is released from its food matrix during digestion and made accessible for absorption into the intestinal mucosa. The food matrix is one of the important factors that relates to the bioaccessibility of carotenoids, and may be changed by heat through cell wall softening. Several studies have shown an increased bioaccessibility of β -carotene in carrots and carrot products by thermal pasteurization (Knockaert et al., 2012; Lemmens et al., 2013). Higher β -carotene bioaccessibility is normally associated with intense thermal processes.

Texture

The texture of processed vegetables is another primary marketable characteristic for customers. Mechanisms that contribute to texture loss during heating of vegetables generally include turgor loss due to the breakdown of cellular membranes, and cell wall degradation and disassembly resulting from enzymatic and nonenzymatic transformations in pectin structure and composition (Anthon et al., 2005; Greve et al., 1994a, b; Peng et al., 2014; Sila et al., 2008). However, not all enzyme-catalyzed reactions to pectin reduce the texture of processed vegetables. For example, the de-esterification of pectin by PME during low temperature blanching contributes to the firming of vegetable tissues. For mild pasteurization in which the processing temperature is lower than 80°C, vegetable tissue softening due to pectin depolymerization by nonenzymatic degradation via β -elimination is negligible due to the relatively high temperature ($T > 80^\circ\text{C}$) and pH levels ($\text{pH} > 4.5$) required for this reaction to take place (Sila et al., 2008).

Texture characteristics of vegetables can be evaluated by sensory and instrumental methods. Sensory evaluation offers the opportunity to obtain a complete analysis of the textural properties of a food as perceived by the human sense, while instrumental measurements are more convenient, less expensive, and tend to provide consistent values when used by different people (Bourne, 1982; Abbott, 2004). Evaluations of vegetable texture in the publications listed in Table 10 were all assessed by instrumental methods, although the specific equipment and method were product-dependent. Most authors applied force-compression tests by using a texture analyzer, only one study by Koo et al. (2008) used a rheometer to measure the texture of soybean sprouts.

Most authors observed a decreased texture in processed vegetables in comparison to the raw materials. A first order reaction model is usually applied to predict the softening of vegetables under pasteurization temperatures (Lau et al., 2000). Processed vegetables may also lose their texture during the storage period, and sometimes this loss in storage may exceed the texture loss during the production process. For example, Koskineniemi et al. (2013) pasteurized sweet potato, red bell pepper and broccoli in a continuous microwave system (3.5 kW) for 4 min at 75°C and held the products in insulating molds for 30 min. They found that the texture of red bell pepper was reduced by half after processing, and was almost completely lost at the end of a 60-day storage period at 30°C. A similar phenomenon of major texture loss occurring during the storage period was observed in the other 2 commodities. The authors stated that one possible reason was the addition of NaCl and citric acid, which may have resulted in tissue softening during the storage time. Besides the presence of these compounds in the media, residual enzyme activity in processed products may also degrade texture through reactions occurring during storage. Rayman and Baysal (2011) investigated residual PME activities and pectin content in pasteurized carrot juice during a 4-month storage at 4°C, but no information on the correlation between the 2 were provided in that study. Schultz et al. (2014) studied the effects of adding PME to pasteurized carrot juice, quantified the juice texture via cloud stability and particle size, and correlated texture with the levels of PME addition during juice

storage (56 days/ 5°C). The authors found that all levels of PME addition (0.15–1.0 U/g) resulted in clarification, and higher amounts had a modest effect in causing more rapid clarification, due to a faster increase in particle size.

Phenolics and antioxidant activity

Phenolics are important phytochemicals that function as bioactive compounds in vegetables. Most researchers have reported phenolics in relation to their antioxidant activity. Effects of thermal pasteurization on the total phenolics in vegetables have been associated with the properties of the food material, package and storage conditions. A reduction in the total phenolic content of pumpkin and carrot juice in pasteurization and post-pasteurization storage at 4°C up to 3 or 4 months was reported by Rayman and Baysal (2011) and Zhou et al. (2014). In contrast, Odriozola-Serrano et al. (2008) did not find significant changes in the total phenolic content between pasteurized and fresh tomato juice (90°C for 30/60 s), and noticed good maintenance of the phenolic compounds during storage. This might be due to the inactivation of the enzymes responsible for their degradation. However, the same authors later reported a decrease in the phenolic concentration of pasteurized tomato juice during storage using the same processing and storage conditions (Odriozola-Serrano et al., 2009). The authors hypothesized that the degradation of phenolic compounds during storage was associated with residual activity of peroxidase, but no enzyme activity assay was carried out to support this assumption. None of the papers that associated phenolic changes in vegetables during storage examined the related enzyme activities (Table 10).

Quercetin is one of the numerous flavonoids commonly found in vegetables. It is known that flavonoids are relatively labile to degradation due to their hydroxyl and ketone groups, and also because of the unsaturated double bonds. Most studies reported a decrease in quercetin content in pasteurized vegetables (Lee and Howard, 1999; Odriozola-Serrano et al., 2009). The changes in quercetin are also associated with the specific processing method. Roldán et al. (2008) evaluated the total quercetin content in pasteurized onion by-products and their frozen products, and found a lower value in all the pasteurized products compared to their corresponding frozen products, but higher quercetin values than the sterilized onion products. With regard to the quercetin changes during storage, stable levels in yellow banana pepper during storage (23°C /4 months) were reported by Lee and Howard (1999), while a decrease in pasteurized tomato juice (23°C /4 months) was observed by Odriozola-Serrano et al. (2009).

Antioxidant capacity is associated with the levels of some bioactive compounds in foods, such as phenolics, vitamin C, and lycopene. Generally, antioxidant activity is reduced in pasteurized vegetables, and this depletion is associated with heat intensity (Rayman and Baysal, 2011). However, Odriozola-Serrano (2008) demonstrated no changes in the antioxidant activity between fresh and pasteurized tomato juice, and explained this by the formation of novel compounds such as products from the Maillard reaction which have antioxidant activity. The authors reported that the antioxidant capacity of pasteurized tomato juice decreased with storage time and attributed this to the losses of vitamin C and lycopene.

Vitamins

Vegetables are great sources of various essential vitamins. Vitamin C (ascorbic acid) is one of the numerous vitamins in vegetables. However, vitamin C is readily changed or broken down in the presence of oxygen and light, and high temperature will accelerate this degradation process. Due to its thermolability, vitamin C in vegetables is often used as an indicator for the loss of other vitamins and thermolabile nutrients in studies that evaluate the influence of thermal processing on food qualities. Thermal degradation of vitamin C in foods is generally reported as a first-order kinetic reaction, both during thermal treatment and storage (Odrizola-Serrano et al., 2008; Torregrosa et al., 2006). Large losses of vitamin C usually occur at the beginning of the thermal treatment, and high temperatures accelerate the degradation process (Elez-Martínez and Martín-Belloso, 2007; Koo et al., 2008; Torregrosa et al., 2006). For example, thermal pasteurization of gazpacho (a cold vegetable soup) at 90°C for 1 min reduced the vitamin C level to 79.2% of its initial value (Elez-Martínez and Martín-Belloso, 2007). Other vitamins, such as vitamin E and D, have also been reported to decrease in vegetable beverages after pasteurization (Barba, 2012).

Most authors have reported reductions in vitamin C during storage following thermal treatment. The lower level depended on storage conditions, such as temperature, oxygen content, light and packaging conditions. The first-order kinetic model has been used to describe the degradation of vitamin C in storage (Odrizola-Serrano et al., 2008). Conceivably, its degradation rate during normal storage is generally lower than during thermal processing. For example, the content of vitamin C decreased by 19.2% in thermally treated pumpkin (85°C for 5 min), and only additional 12.1% loss after 2 months storage at 4°C (Zhou et al., 2014). Low storage temperatures usually slow down ascorbic acid degradation rate (Torregrosa et al., 2006). However, Koo et al. (2008) reported that a decreased level of ascorbic acid in cook-chilled packaged sprouts stored at 3°C was much like those stored at 10°C, which may be due to a relatively low level of this nutrient after thermal treatment.

Other components

Changes in other nutrients or relevant quality aspects, such as sugars, dry matter content, fatty acids, isothiocyanates and furfural have been reported in vegetables during thermal treatments and storage (Vervoort et al., 2012; Zhou et al., 2014), this information is not included in Table 10. Vervoort et al. (2012) reported a significant reduction in dry matter content, furfural, and sugar concentration (glucose, fructose and sucrose) of carrot pieces during pasteurization, and found that increasing the processing intensity from a level of $F_{70^{\circ}\text{C}} = 2$ min to $F_{90^{\circ}\text{C}} = 10$ min did not cause any further significant changes in those values. However, the study by Zhou et al. (2014) showed that pasteurization did not significantly change the sugar content (sucrose, glucose and fructose) in pumpkin after thermal treatment (85°C/ 5 min) and during storage (60 days/4°C).

Storage and shelf-life of pasteurized vegetables

Storage is an important component for the entire food chain to ensure the safety and quality of foods from the field to the table. Many storage conditions, such as storage temperature, relative humidity, air circulation, ventilation, packaging, stacking, volatile substances, and hygiene may affect the quality of processed products. For pasteurized foods, refrigerated storage is normally required to extend the shelf-life of the products. Refrigeration slows down the growth of spoilage microorganisms and the rate of chemical changes in foods, therefore it reduces the food deterioration rate. Table 10 summarizes the published papers on pasteurized vegetables related to storage and enzyme studies. Few studies investigated the residual enzyme activity after processing. It is also notable that storage temperatures and time in the works cited varied from product to product. For most of the pasteurized vegetables, a storage temperature of 3–5°C was used. For some pickled, high acid vegetables, the storage temperature was much higher (23–30°C) because of the low pH of the products. The storage times reported in these publications varied from 21 days to 5 months, depending on the property of the products and the storage temperature. Aside from pH as mentioned above, other factors such as reduced water activity, preservatives, or a combination thereof can be used to control pathogen growth or spoilage during storage and affect the storage life of the products.

With regard to the shelf-life of pasteurized ready-to-eat foods in general and vegetables in particular, there does not appear to be a universally accepted standard for all products. Torregrosa et al. (2006) calculated the shelf-life of pasteurized orange-carrot juice as the time taken for the ascorbic acid concentration to reduce by 50% (Table 10). Most authors do not provide reasons for the storage conditions they selected for their products. Shelf life is defined as “the period of time for which a product remains safe and meets its quality specifications under expected storage and use conditions” (ECFF, 2006). Based on the ECFF (2006), the manufacturer is responsible for determining the shelf life and must take into account microbiological safety and stability, physical characteristics and organoleptic quality. Microbiological safety and stability should always be a priority for determining the shelf life when the acceptable shelf life for either physical condition or organoleptic quality exceeds that for microbial safety. The product shelf life is influenced by a number of factors, including raw material quality, product formulation (pH, a_w), hygiene during manufacturing, scheduled heat or other preservation treatments, cooling methods applied to products, type of package, storage temperature and relevant hurdles (CAC, 1999). When determining the shelf life of products, the Codex Alimentarius Commission (CAC) suggests taking into consideration the potential for temperature abuse which may occur during manufacture, storage, distribution, sale, and handling by the consumer. For example, fluid milk is most often held at marginal refrigeration temperatures of 6.1–7.2°C (43–45°F) instead of the ideal holding temperatures ($\leq 3.3^{\circ}\text{C}$) to determine the potential shelf-life (Murphy, 2009). The authors believed these marginal refrigeration temperatures allow defects and sanitation deficiencies to become more evident. The FDA requires control measures for storage under normal and moderate abuse

conditions (FDA, 2011b), for example, the recommended incubation temperature for inoculated pack studies with nonproteolytic *C. botulinum* is 10°C (50°F) (NACMCF, 1992). Therefore, for a food producer determining the safe shelf-life for pasteurized ready-to-eat foods and vegetables, the following information needs to be collected based on ECFE recommendations:

- (1) Review relevant scientific information containing the characteristics of pathogens;
- (2) Use predictive modeling programs (e.g., ComBase, USFA Pathogen Modeling Program or Growth Predictor) to estimate the growth of pathogens under the storage conditions;
- (3) Conduct a challenge test with the relevant pathogens where predictive modeling on its own does not give sufficient confidence to set a safe shelf life;
- (4) Collect historical data for similar products;
- (5) Conduct storage trials, either by storing products at predetermined temperatures during specific time periods considering actual chill chain performance under HACCP or testing the product at minimum 3 time points for the relevant indicator and spoilage microorganisms as well as pathogens identified by HACCP.

The information above focuses on a safe shelf life. Desired quality should also be considered when determining the shelf-life of pasteurized products. This information also provides insights for researchers to conduct storage study for pasteurized products.

Conclusions and future trends

To design a thermal pasteurization process for ready-to-eat meals including vegetable products that could meet the regulations and guidelines of governmental food agencies, it is of paramount importance to fully understand regulatory requirements and select the pathogens of concern that could survive the process and storage conditions. Determination of the target pathogens for a pasteurization process according to the U.S. regulations is product and process dependent; there is no “one size fits all” approach. Despite the differences in the specific requirements for target pathogens for pasteurized foods in the United States and Europe, safety storage conditions throughout product shelf-life are common concerns. Consideration of moderate temperature abuse during storage and distribution should be included when determining the target pathogens for the process design.

Viral pathogens are an emerging concern for foods, and progress has been made in characterizing the environmental resistance (especially thermal resistance) of some human norovirus surrogates. Based on our current knowledge, norovirus or HAV may be able to exist in some low temperature processed foods. Therefore, to develop a pasteurization process, it is suggested that the presence of viral pathogens be taken into consideration in addition to the common pathogens, particularly in fruit and vegetables, shellfish, deli meats and ready-to-eat foods.

After addressing the safety concern for a pasteurized product, attention should be given to the product quality changes due to the storage and distribution of the food. Thermal kinetic

data of quality-associated enzymes and quality parameters in vegetables are available in the literature, and can be used for the pasteurization process optimization. To develop a pasteurized vegetable product using a new technology, storage trials should be carried out at normal distribution and storage conditions to verify the pasteurization process and ensure safe and quality-acceptable products within their maximum shelf-life.

Acknowledgments

The authors thank the Chinese Scholarship Council for providing a scholarship to Jing Peng for her PhD Studies at Washington State University prior to her appointment at Nanjing Agricultural University.

Funding

This research was funded by the U.S. Agriculture and Food Research Initiative of the USDA National Institute of Food and Agriculture, Grant Number #2011–68003–20096, & #2016–68003–20096, and partially by the Fundamental Research Funds for the Central Universities of China (KYZ201655).

References

- Aamir, M., Ovissipour, M., Rasco, B., Tang, J. and Sablani, S. (2014). Seasonality of thermal kinetics of color changes in whole spinach (*Spinacea oleracea*) leaves under pasteurization conditions. *Int. J. Food Prop.* **17**:2012–2024.
- Aamir, M., Ovissipour, M., Sablani, S. S. and Rasco, B. (2013). Predicting the quality of pasteurized vegetables using kinetic models: A review. *Int. J. Food Sci.* DOI: 10.1155/2013/271271.
- Abbott, J. A. (2004). Sensory and instrumental measurement of texture of fruits and vegetables. *Hortscience* **39**:830.
- Agüero, M. V., Ansorena, M. R., Roura, S. I. and del Valle, C. E. (2008). Thermal inactivation of peroxidase during blanching of butternut squash. *LWT - Food Sci. Technol.* **41**:401–407.
- Ahmed, N. M., Conner, D. E. and Huffman, D. L. (1995). Heat resistance of *Escherichia coli* O157:H7 in meat and poultry as affected by product composition. *J. Food Sci.* **60**:606–610.
- Andrés-Bello, A., Barreto-Palacios, V., García-Segovia, P., Mir-Bel, J. and Martínez, Monzó. (2013). Effect of pH on color and texture of food products. *Food Eng. Rev.* **5**:158–170.
- Anese, M. and Sovrano, S. (2006). Kinetics of thermal inactivation of tomato lipoxygenase. *Food Chem.* **95**:131–137.
- Anthon, G. and Barrett, D. M. (2002). Kinetic parameters for the thermal inactivation of quality-related enzymes in carrots and potatoes. *J. Agric. Food Chem.* **50**:4119–4125.
- Anthon, G. E. and Barrett, D. M. (2003). Thermal inactivation of lipoxygenase and hydroperoxytrieneic acid lyase in tomatoes. *Food Chem.* **81**:275–279.
- Anthon, G. and Barrett, D. M. (2006). Characterization of the temperature activation of pectin methylesterase in green beans and tomatoes. *J. Agric. Food Chem.* **54**:204–211.
- Anthon, G. E., Blot, L. and Barrett, D. M. (2005). Improved firmness in calcified diced tomatoes by temperature activation of pectin methylesterase. *J. Food Sci.* **70**:C342–C347.
- Audits International/FDA. (1999). U.S. cold temperature evaluation. Available at: http://foodrisk.org/default/assets/File/Audits-FDA_temp_study.pdf. Accessed 9 August 2015.
- Barba, F. J., Esteve, M. J. and Frigola, A. (2012). Impact of high-pressure processing on vitamin E (α -, γ -, and δ -Tocopherol), vitamin D (cholecalciferol and ergocalciferol) and fatty acid profiles in liquid foods. *J. Agric. Food Chem.* **60**:3763–3768.
- Baert, L., Uyttendaele, M., Van Coillie, E. and Debevere, J. (2008). The reduction of murine norovirus 1, B. fragilis HSP40 infecting phage B40–8 and E. coli after a mild thermal pasteurization process of raspberry puree. *Food Microbiol.* **25**:871–874.

- Balogh, T., Smout, C., Ly-Nguyen, B., Loey, A. V. and Hendrickx, M. E. (2004). Thermal and high-pressure inactivation kinetics of carrot pectinmethylesterase: From model system to real foods. *Innov. Food Sci. Emerg. Technol.* **5**:429–436.
- Barrett, D. M. (1994). Vegetable blanching—How long is long enough? *Am. Vegetable Grower* **42**(1):46–47.
- Bidawid, S., Farber, J. M., Sattar, S. A. and Hayward, S. (2000). Heat inactivation of hepatitis A virus in dairy foods. *J. Food Prot.* **63**:522–528.
- Bolton, D. J., McMahon, C. M., Doherty, A. M., Sheridan, J. J., McDowell, D. A., Blair, L. S. and Harrington, D. (2000). Thermal inactivation of *Listeria monocytogenes* and *Y. enterocolitica* in minced beef under laboratory conditions and in sous-vide prepared minced and solid beef cooked in a commercial retort. *J. Appl. Microbiol.* **88**:626–632.
- Bourne, M. C. (1982). *Food Texture and Viscosity: Concept and Measurement*. Academic Press, New York.
- Bozkurt, H., D'Souza, D. H. and Davidson, P. M. (2013). Determination of the thermal inactivation kinetics of the human norovirus surrogates, murine norovirus and feline calicivirus. *J. Food Prot.* **76**:79–84.
- Bozkurt, H., D'Souza, D. H. and Davidson, P. M. (2014a). Determination of thermal inactivation kinetics of hepatitis A virus in blue mussel (*Mytilus edulis*) homogenate. *Appl. Environ. Microbiol.* **80**:3191–3197.
- Bozkurt, H., D'Souza, D. H. and Davidson, P. M. (2014b). A comparison of the thermal inactivation kinetics of human norovirus surrogates and hepatitis A virus in buffered cell culture medium. *Food Microbiol.* **80**:212–217.
- Bozkurt, H., D'Souza, D. H. and Davidson, P. M. (2014c). Thermal inactivation of human norovirus surrogates in spinach and measurement of its uncertainty. *J. Food Prot.* **77**:276–283.
- Bozkurt, H., Leiser, S., Davidson, P. M. and D'Souza, D. H. (2014d). Thermal inactivation kinetic modeling of human norovirus surrogates in blue mussel (*Mytilus edulis*) homogenate. *Int. J. Food Microbiol.* **172**:130–136.
- Bozkurt, H., D'Souza, D. H. and Davidson, P. M. (2015a). Thermal inactivation kinetics of human norovirus surrogates and hepatitis A virus in turkey deli meat. *Appl. Environ. Microbiol.* **81**:4850–4859.
- Bozkurt, H., D'Souza, D. H. and Davidson, P. M. (2015b). Thermal inactivation of hepatitis A virus in homogenized clams (*Mercenaria mercenaria*). *J. Appl. Microbiol.* **119**:834–844.
- Bozkurt, H., Ye, X., Harte, F., D'Souza, D. H. and Davidson, P. M. (2015c). Thermal inactivation kinetics of hepatitis A virus in spinach. *Int. J. Food Microbiol.* **193**:147–151.
- Buckow, R., Isbarn, S., Knorr, D., Heinz, V. and Lehmacher, A. (2008). Predictive model for inactivation of feline calicivirus, a norovirus surrogate, by heat and high hydrostatic pressure. *Appl. Environ. Microbiol.* **74**:1030–1038.
- Buduamoako, E., Toora, S., Walton, C., Ablett, R. F. and Smith, J. (1992). Thermal death times for *Listeria monocytogenes* in lobster meat. *J. Food Prot.* **55**:211–213.
- Cannon, J. L., Papafragkou, E., Park, G. W., Osborne, J., Jaykus, L. J. and Vinjé, J. (2006). Surrogates for the study of norovirus stability and inactivation in the environment: a comparison of murine norovirus and feline calicivirus. *J. Food Prot.* **69**:2761–2765.
- CDC. (2006). Botulism associated with commercial carrot juice—Georgia and Florida, September 2006. Morbidity and Mortality Weekly Report (MMWR). **55**(Dispatch):1–2. Centers for Disease Control. Available at <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm55d106a1.htm>. Accessed October 6, 2006.
- CDC. (2011). CDC estimates of foodborne illness in the United States. Centers for Disease Control and Prevention. Available at: <http://www.cdc.gov/foodborneburden/2011-foodborne-estimates.html#illness>
- Cerf, O. and Condrón, R. (2006). *Coxiella burnetii* and milk pasteurization: an early application of the precautionary principle? *Epidemiol. Infect.* **134**:946–951.
- Code of Federal Regulations. (2009). 21 CFR 118. *Prevention of Salmonella Enteritidis in Shell Eggs during Production, Storage, and Transportation*. U.S. Government Printing Office, Washington, D.C. Federal Register. **74**(130):33030–33101.
- Code of Federal Regulations. (2011). 21 CFR 120.24 U.S. Government Printing Office, Washington, D.C..
- Code of Federal Regulations. (2012a). 9 CFR 590. *Inspection of Egg and Egg Products*. U.S. Government Printing Office, Washington, D.C..
- Code of Federal Regulations. (2012b). 9 CFR 590:570. U.S. Government Printing Office, Washington, D.C..
- Code of Federal Regulations. (2012c). 9 CFR 590:575. U.S. Government Printing Office, Washington, D.C..
- Codex Alimentarius Commission. (1999). Codex code of hygienic practice for refrigerated packaged foods with extended shelf life, CAC/RCP 46.
- Croci, L., Ciccozzi, M., De Medici, D., Di Pasquale, S., Fiore, A., Mele, A. and Toti, L. (1999). Inactivation of hepatitis A virus in heat-treated mussels. *J. Appl. Microbiol.* **87**:884–888.
- Croci, L., De Medici, D., Di Pasquale, S. and Toti, L. (2005). Resistance of hepatitis A virus in mussels subjected to different domestic cooking. *Int. J. Food Microbiol.* **105**:139–144.
- Croci, L., Suffredini, E., Di Pasquale, S. and Cozzi, L. (2012). Detection of *Norovirus* and *Feline Calicivirus* in spiked mollusks subjected to heat treatments. *Food Contr.* **25**:17–22.
- CSIRO Food and Nutritional Sciences. (2010). *Make it Safe: A Guide to Food Safety*. CSIRO Publishing, Collingwood.
- De-Angelis, M., Di-Cagno, R., Huet, C., Crecchio, C., Fox, P. F. and Gobetti, M. (2004). Heat shock response in *Lactobacillus plantarum*. *Appl. Environ. Microbiol.* **70**:1336–1346.
- Deboosere, N., Pinon, A., Delobel, A., Temmam, S., Morin, T., Merle, G., Blaise-Boisseau, S., Perelle, S. and Vialette, M. (2010). A predictive microbiology approach for thermal inactivation of hepatitis A virus in acidified berries. *Food Microbiol.* **27**:962–967.
- Deboosere, N., Legeay, O., Caudrelier, Y. and Lange, M. (2004). Modelling effect of physical and chemical parameters on heat inactivation kinetics of hepatitis A virus in a fruit model system. *Int. J. Food Microbiol.* **93**:73–85.
- Dolin, R., Blacklow, N., DuPont, H., Buscho, R., Wyatt, R., Kasel, J., Homick, R. and Chanock, R. (1972). Biological properties of Norwalk agent of acute infectious nonbacterial gastroenteritis. *P. Soc. Exp. Biol. Med.* **140**:578–583.
- Doultree, J. C., Druce, J. D., Birch, C. J., Bowden, D. S. and Marshall, J. A. (1999). Inactivation of feline calicivirus, a Norwalk virus surrogate. *J. Hosp. Infect.* **41**:51–57.
- Downing, D. L. (1996). *A Complete Course in Canning and Related Processes* 13th ed-Book II: Microbiology, Packaging, HACCP & Ingredients. Woodhead Publishing, Cambridge, UK.
- Duizer, E., Bijkerk, P., Rockx, B., de Groot, A., Twisk, F. and Koopmans, M. (2004). Inactivation of caliciviruses. *Appl. Environ. Microbiol.* **70**:4538–4543.
- European Food Safety Authority (EFSA). (2011). Scientific opinion on an update on the present knowledge on the occurrence and control of foodborne viruses. Available at <http://www.efsa.europa.eu/fr/efsajournal/doc/2190.pdf>. Accessed 9 August 2015.
- European Chilled Food Federation (ECFF). (2006). *Recommendations for the production of prepacked chilled food*, 2nd ed. Available at http://www.ecff.net/images/ECFF_Recommendations_2nd_ed_18_12_06.pdf. Accessed 9 August 2015.
- Elez-Martínez, P. and Martín-Belloso, O. (2007). Effects of high intensity pulsed electric field processing conditions on vitamin C and antioxidant capacity of orange juice and gazpacho, a cold vegetable soup. *Food Chem.* **102**:201–209.
- Enright, J. B., Sadler, W. W. and Thomas, R. C. (1957). Thermal inactivation of *Coxiella burnetii* and its relation to pasteurization of milk. *Public Health Monogr.* No. 47.
- Enright, J. B. (1961). The pasteurization of cream, chocolate milk and ice cream mixes containing the organism of Q fever. *J. Milk Food Technol.* **24**:351–355.
- Ercan, S. S. and Soysal, Cigdem. (2011). Effect of ultrasound and temperature on tomato peroxidase. *Ultrason. Sonochem.* **18**:689–695.
- Espach-Barroso, A., Loey, A. V., Hendrickx, M. and Martín-Belloso, O. (2006). Inactivation of plant pectin methylesterase by thermal or high intensity pulsed electric field treatments. *Innov. Food Sci. Emerg. Technol.* **7**:40–48.
- European Commission (EC). (2004). Corrigendum to Regulation (EC) No 852/2004 of the European Parliament and of the Council of 29 April

- 2004 on the hygiene of foodstuffs. *Off. J. Eur. Union*. **L139**, 30 April 2004.
- Fachin, D., Van Loey, A. M., Ly Nguyen, B., Verlent, I. and Hendrickx, M. E. (2003). Inactivation kinetics of polygalacturonase in tomato juice. *Innov. Food Sci. Emerg. Technol.* **4**:135–142.
- FDA. (2004). Guidance for Industry: Juice HACCP Hazards and Controls Guidance 1st ed; final guidance. U.S. Food and Drug Administration, Department of Health and Human Service. Available at <http://www.fda.gov/Food/GuidanceRegulation/GuidanceDocumentsRegulatoryInformation/Juice/ucm072557.htm>. Accessed 9 August 2015.
- FDA. (2011a). Grade “A” Pasteurized Milk Ordinance. U.S. Food and Drug Administration, Department of Health and Human Services. Available at <http://www.fda.gov/downloads/Food/GuidanceRegulation/UCM291757.pdf>. Accessed 9 August 2015.
- FDA. (2011b). Fish and fisheries products hazards and control guidance, 4th ed. U.S. Food and Drug Administration, Department of Health and Human Services. Available at <http://www.fda.gov/downloads/food/guidanceregulation/ucm251970.pdf>. Accessed 9 August 2015.
- FFDCA. (1938). *Federal Food Drug and Cosmetic Act (June 25, 1938)*. Government Printing Office, Washington D.C. Available at <http://www.gpo.gov/fdsys/pkg/USCODE-2010-title21/pdf/USCODE-2010-title21-chap9.pdf>. Accessed 9 August 2015.
- Fernández, A., Ocio, M. J., Fernández, P. S. and Martínez, A. (2001). Effect of heat activation and inactivation conditions on germination and thermal resistance parameters of *Bacillus cereus* spores. *Int. J. Food Microbiol.* **63**:257–264.
- FSMA. (2011). *Food Safety Modernization Act. PL 111–353 (Jan. 4, 2011)*. Government Printing Office, Washington D.C. Available at <http://www.gpo.gov/fdsys/pkg/PLAW-111publ353/pdf/PLAW-111publ353.pdf>. Accessed 9 August 2015.
- Franz, C. M. A. P. and vonHoly, A. (1996). Thermotolerance of meat spoilage lactic acid bacteria and their inactivation in vacuum packaged vienna sausages. *Int. J. Food Microbiol.* **29**:59–73.
- Gaze, J. E. and Brown, G. D. (1990). Determination of the heat resistance of a strain of non-proteolytic *Clostridium botulinum* type B and a strain of type E, heated in cod and carrot homogenate over the temperature range 70 to 92°C. Campden Food and Drink Research Association Technical Memorandum N. 592. Chipping Campden, UK.
- Gibson, K. E. and Schwab, K. J. (2011). Thermal inactivation of human norovirus surrogates. *Food Environ. Virol.* **3**:74–77.
- Gonçalves, E. M., Pinheiro, J., Abreu, M., Brandão, T. R. S. and Silva, C. L. M. (2010). Carrot (*Daucus carota* L.) peroxidase inactivation, phenolic content and physical changes kinetics due to blanching. *J. Food Eng.* **97**:574–581.
- Greve, L. C., McArdle, R. N., Gohlke, J. R. and Labavitch, J. M. (1994a). Impact of heating on carrot firmness: Changes in cell wall components. *J. Agric. Food Chem.* **42**:2900–2906.
- Greve, L. C., Shackel, K. A., Ahmadi, H., McArdle, R. N., Gohlke, J. R. and Labavitch, J. M. (1994b). Impact of heating on carrot firmness: Contribution of cellular turgor. *J. Agric. Food Chem.* **42**:2896–2899.
- Guivarçh, Y., Sila, D., Duvetter, T., Van Loey, A. and Hendrickx, M. (2003). Influence of sugars and polyols on the thermal stability of purified tomato and cucumber pectinmethylesterases: A basis for TTI development. *Enzyme Microb. Tech.* **33**:544–555.
- Harlow, J., Oudit, D., Hughes, A. and Mattison, K. (2011). Heat inactivation of hepatitis A virus in shellfish using steam. *Food Environ. Virol.* **3**:31–34.
- Hernández-Carrión, M., Hernando, I. and Quiles, A. (2014). High hydrostatic pressure treatment as an alternative to pasteurization to maintain bioactive compound content and texture in red sweet pepper. *Innov. Food Sci. Emerg. Technol.* **26**:76–85.
- Hewitt, J., Rivera-Aban, M. and Greening, G. E. (2009). Evaluation of murine norovirus as a surrogate for human norovirus and hepatitis A virus in heat inactivation studies. *J. Appl. Microbiol.* **107**:65–71.
- Hirneisen, K. A., Black, E. P., Cascarino, J. L., Fino, V. R., Hoover, D. G. and Kniel, K. E. (2010). Viral inactivation in foods: A review of traditional and novel food-processing technologies. *Compr. Rev. Food Sci. F.* **9**:3–20.
- Horm, K. M. and D’Souza, D. H. (2011). Survival of human norovirus surrogates in milk, orange, and pomegranate juice, and juice blends at refrigeration (4°C). *Food Microbiol.* **28**:1054–1061.
- Howard, L. R., Burma, P. and Wagner, A. B. (1997). Firmness and cell wall characteristics of pasteurized Jalapeño pepper rings as affected by pre-heating and storage. *J. Food Sci.* **62**:89–112.
- Huang, I. P. D., Yousef, A. E., Marth, E. H. and Matthews, M. E. (1992). Thermal inactivation of *Listeria monocytogenes* in chicken gravy. *J. Food Prot.* **55**:492–496.
- Indrawati Van Loey, A. M., Ludikhuyze, L. R. and Hendrickx, M. E. (1999). Single, combined, or sequential action of pressure and temperature on lipoxygenase in green beans (*Phaseolus vulgaris* L.): A kinetic inactivation study. *Biotechnol. Prog.* **15**:273–277.
- Indrawati, I., Van Loey, A., Ludikhuyze, L. and Hendrickx, M. (2001). Pressure-temperature inactivation of lipoxygenase in green peas (*Pisum sativum*): A kinetic study. *J. Food Sci.* **66**:686–693.
- Jin, T., Zhang, H., Boyd, G. and Tang, J. (2008). Thermal resistance of *Salmonella Enteritidis* and *Escherichia coli* K12 in liquid egg determined by thermal-death-time disks. *J. Food Eng.* **84**:608–614.
- Johne, R., Trojnar, E., Filter, M. and Hofmann, J. (2016). Thermal stability of hepatitis E virus estimated by a cell culture method. *Appl. Environ. Microbiol.* **82**:4225–4231.
- Juneja, V. K. (2007). Thermal inactivation of *Salmonella spp.* in ground chicken breast or thigh meat. *Int. J. Food Sci. Technol.* **42**:1443–1448.
- Juneja, V. K. and Marmer, B. S. (1999). Lethality of heat to *Escherichia coli* O157:H7: D- and z-value determinations in turkey, lamb and pork. *Food Res. Int.* **32**:23–28.
- Juneja, V. K., Snyder, O. P. and Marmer, B. S. (1997). Thermal destruction of *Escherichia coli* O157:H7 in beef and chicken: determination of D- and z-values. *Int. J. Food Microbiol.* **35**:231–237.
- Jung, Y. S. and Beuchat, L. R. (2000). Sensitivity of multidrug-resistant *Salmonella typhimurium* DT104 to organic acids and thermal inactivation in liquid egg products. *Food Microbiol.* **17**:63–71.
- Keswani, J. and Frank, J. F. (1998). Thermal inactivation of *Mycobacterium paratuberculosis* in milk. *J. Food Prot.* **61**:974–978.
- Knockaert, G., Lemmens, L., Van Buggenhout, S., Hendrickx, M. and Van Loey, A. (2012). Changes in β -carotene bioaccessibility and concentration during processing of carrot puree. *Food Chem.* **133**:60–67.
- Koo, K. M., Kim, H. W., Lee, D. S., Lyu, E. S. and Paik, H. D. (2008). Quality changes during storage of cook-chilled soybean sprouts. *Food Sci. Biotechnol.* **17**:540–546.
- Koopmans, M. and Duizer, E. (2004). Foodborne viruses: an emerging problem. *Int. J. Food Microbiol.* **90**:23–41.
- Koskiniemi, G. B., Truong, V. D., McFeeters, R. F. and Simunovic, J. (2013). Quality evaluation of packaged acidified vegetables subjected to continuous microwave pasteurization. *LWT-Food Sci. Technol.* **54**:157–164.
- Lau, M. H. and Tang, J. (2002). Pasteurization of pickled asparagus using 915 MHz microwaves. *J. Food Eng.* **51**:283–290.
- Lau, M. H., Tang, J. and Swanson, B. G. (2000). Kinetics of textural and color changes in green asparagus during thermal treatments. *J. Food Eng.* **45**:231–236.
- Lee, Y. and Howard, L. (1999). Firmness and phytochemical losses in pasteurized yellow banana peppers (*Capsicum annuum*) as affected by calcium chloride and storage. *J. Agric. Food Chem.* **47**:700–703.
- Lemmens, L., Colle, I., Knockaert, G., Van Buggenhout, S., Van Loey, A. and Hendrickx, M. (2013). Influence of pilot scale in pack pasteurization and sterilization treatments on nutritional and textural characteristics of carrot pieces. *Food Res. Int.* **50**:525–533.
- Ling, B., Tang, J., Kong, F., Mitcham, E. J. and Wang, S. (2015). Kinetics of food quality changes during thermal processing: A review. *Food Bioprocess Technol.* **8**:343–358.
- Ly-Nguyen, B., Loey, A. V., Fachin, D., Verlent, I., Wati, I. and Hendrickx, M. (2002). Partial purification, characterization and thermal and high-pressure inactivation of pectin methylesterase from carrots (*Daucus carota* L.). *J. Agric. Food Chem.* **50**:5437–5444.
- Ly-Nguyen Loey, A. V., Fachin, D., Verlent, I., Wati, I. and Hendrickx, M. (2003). Mild-heat and high pressure inactivation of carrot pectin methylesterase: A kinetic study. *J. Food Sci.* **4**:1377–1383.
- Loong, M. N. and Goh, H. K. (2004). Colour degradation of acidified vegetable juice. *Int. J. Food Sci. Technol.* **39**:437–441.

- Meanwell, L. J. (1927). An investigation into the effect of pasteurization on the bovine tubercle *Bacillus* in naturally infected tuberculous milk. *J. Hyg.* **26**:392–402.
- Miettinen, H., Arvola, A. and Wirtanen, G. (2005). Pasteurization of rainbow trout roe: *Listeria monocytogenes* and sensory analyses. *J. Food Prot.* **68**:1641–1647.
- Morales-Blancas, E. F., Chandia, V. E. and Cisneros-Zevallos, L. (2002). Thermal inactivation kinetics of peroxidase and lipoxigenase from broccoli, green asparagus and carrot. *Food Chem. Toxicol.* **67**:146–154.
- Murphy, S. C. (2009). Shelf-life of fluid milk products—microbial spoilage—The evaluation of shelf-life. *Dairy Foods Science Notes*. Cornell Univ. Available at <http://foodscience.cornell.edu/cals/foodsci/extension/upload/CU-DFScience-Notes-Bacteria-Milk-Shelf-Life-Evaluation-06-09.pdf>. Accessed 9 August 2015.
- Murphy, R. Y., Beard, B. L., Martin, E. M., Duncan, L. K. and Marcy, J. A. (2004a). Comparative study of thermal inactivation of *Escherichia coli* O157:H7, *Salmonella*, and *Listeria monocytogenes* in ground pork. *J. Food Sci.* **69**:97–101.
- Murphy, R. Y., Davidson, M. A. and Marcy, J. A. (2004b). Process lethality prediction for *Escherichia coli* O157:H7 in raw franks during cooking and fully cooked franks during post-cook pasteurization. *J. Food Sci.* **69**:112–116.
- National Advisory Committee on Microbiological Criteria for Foods (NACMCF). (1992). Vacuum or modified atmosphere packaging for refrigerated raw fishery products. Adopted March 20, 1992. Accessed at http://www.fsis.usda.gov/OPHS/nacmcf/past/map_fishery.htm. Accessed 9 August 2015.
- National Advisory Committee on Microbiological Criteria for Foods (NACMCF). (2006). Requisite scientific parameters for establishing the equivalence of alternative methods of pasteurization. *J. Food Prot.* **69**:1190–1216.
- Odrozola-Serrano, I., Soliva-Fortuny, R. and Martín-Belloso, O. (2008). Changes of health-related compounds throughout cold storage of tomato juice stabilized by thermal or high intensity pulsed electric field treatments. *Innov. Food Sci. Emerg. Technol.* **9**:272–279.
- Odrozola-Serrano, I., Soliva-Fortuny, R., Hernández-Jover, T. and Martín-Belloso, O. (2009). Carotenoid and phenolic profile of tomato juices processed by high intensity pulsed electric field compared with conventional thermal treatments. *Food Chem.* **112**:258–266.
- Oteiza, J. M., Giannuzzi, L. and Califano, A. N. (2003). Thermal inactivation of *Escherichia coli* O157:H7 and *Escherichia coli* isolated from morcilla as affected by composition of the product. *Food Res. Int.* **36**:703–712.
- Pearce, L. E., Truong, H. T., Crawford, R. A., Yates, G. F. and Cavaignac Sde Lisle, G. W. (2001). Effect of turbulent-flow pasteurization on survival of *Mycobacterium avium* subsp. paratuberculosis added to raw milk. *Appl. Environ. Microbiol.* **67**:3964–3969.
- Peng, J., Tang, J., Barrett, D. M., Sablani, S. S. and Powers, J. (2014). Kinetics of carrot texture degradation under pasteurization conditions. *J. Food Eng.* **122**:84–91.
- Rayman, A. and Baysal, T. (2011). Yield and quality effects of electroporation and microwave application on carrot juice production and storage. *J. Food Sci.* **76**:C598–C605.
- Rejano, L., Sanchez, A. H., De Castro, A. and Montano, A. (1997). Chemical characteristics and storage stability of pickled garlic prepared using different processes. *J. Food Sci.* **62**:1120–1123.
- Rodriguez-Amaya, D. B. and Kimura, M. (2004). *HarvestPlus Handbook for Carotenoid Analysis*. HarvestPlus, Washington D.C..
- Roldán, E., Sánchez-Moreno, C., de Ancos, B. and Cano, M. P. (2008). Characterization of onion (*Allium cepa* L.) by-products as food ingredients with antioxidant and antibrowning properties. *Food Chem.* **108**:907–916.
- Sair, A. I., D'Souza, D. H. and Jaykus, L. A. (2002). Human enteric viruses as causes of foodborne disease. *Compr. Rev. Food Sci. F.* **1**:73–89.
- Scheerens, J. C. (2001). Phytochemicals and the consumer: Factors affecting fruit and vegetable consumption and the potential for increasing small fruit in the diet. *HortTechnology* **11**:547–556.
- Schultz, A. K., Anthon, G. E., Dungan, S. R. and Barrett, D. M. (2014). Effect of pectin methylesterase on carrot (*Daucus carota*) juice cloud stability. *J. Agric. Food Chem.* **62**:1111–1118.
- Schuman, J. D. and Sheldon, B. W. (1997). Thermal resistance of *Salmonella* spp. and *Listeria monocytogenes* in liquid egg yolk and egg white. *J. Food Prot.* **60**:634–638.
- Schuman, J. D., Sheldon, B. W. and Foegeding, P. M. (1997). Thermal resistance of *Aeromonas hydrophila* in liquid whole egg. *J. Food Prot.* **60**:231–236.
- Selby, T. L., Berzins, A., Gerrard, D. E., Corvalan, C. M., Grant, A. L. and Linton, R. H. (2006). Microbial heat resistance of *Listeria monocytogenes* and the impact on ready-to-eat meat quality after post-package pasteurization. *Meat Sci.* **74**:425–434.
- Seo, K., Lee, J. E., Lim, M. Y., Ko, G. (2012). Effect of temperature, pH, and NaCl on the inactivation kinetics of murine norovirus. *J. Food Prot.* **75**:533–540.
- Shi, J., Maguer, M. L., Bryan, M. and Kakuda, Y. (2003). Kinetics of lycopene degradation in tomato puree by heat and light irradiation. *J. Food Process Eng.* **25**:485–498.
- Sila, D. N., Duvetter, T., Roeck, A. D., Verlent, I., Smout, C., Moates, G. K., Hills, B. P., Waldron, K. K., Hendrickx, M. and Loey, A. V. (2008). Texture changes of processed fruits and vegetables: Potential use of high-pressure processing. *Trends Food Sci. Technol.* **19**:309–319.
- Silva, F. V. M. and Gibbs, P. A. (2010). Non-proteolytic *Clostridium Botulinum* spores in low-acid cold-distributed foods and design of pasteurization processes. *Trends Food Sci. Technol.* **21**:95–105.
- Silva, F. V. M. and Gibbs, P. A. (2012). Thermal pasteurization requirements for the inactivation of *Salmonella* in foods. *Food Res. Int.* **45**:695–699.
- Silva, F. V. M., Gibbs, P. A., Nunez, H., Almonacid, S. and Simpson, R. (2014). *Thermal Processes: Pasteurization*. Encyclopedia of Food Microbiology, 577–595. 2nd edition. Academic Press, San Diego.
- Stanley, D. W., Bourne, M. C., Stone, A. P. and Wismer, W. V. (1995). Low temperature blanching effects on chemistry firmness and structure of canned green beans and carrots. *J. Food Sci.* **60**:327–333.
- Steinfeld, J. I., Francisco, J. S. and Hase, W. L. (1998). *Chemical Kinetics and Dynamics*. 2nd ed. Prentice Hall, Upper Saddle River, NJ.
- Teixeira, A. A. (2014). Thermal food preservation technology (pasteurization, sterilization, canning, and blanching). in *Conventional and Advanced Food Processing Technologies* (ed S. Bhattacharya). John Wiley & Sons, Ltd, Chichester, UK.
- Terefe, N. S., Buckow, R. and Versteeg, C. (2014). Quality-related enzymes in fruit and vegetable products: Effects of novel food processing technologies, part I: High pressure processing. *Crit. Rev. Food Sci. Nutr.* **54** (1):24–63.
- Thomas, C. T., White, J. C. and Longree, K. (1966). Thermal resistance of *Salmonellae* and *Staphylococci* in foods. *Appl. Microbiol.* **14**:815–820.
- Toora, S., Buduamoako, E., Ablett, R. F. and Smith, J. (1992). Effect of high-temperature short-time pasteurization, freezing and thawing and constant freezing, on the survival of *Yersinia enterocolitica* in milk. *J. Food Prot.* **55**:803–805.
- Torregrosa, F., Esteve, M. J., Frígola, A. and Cortés, C. (2006). Ascorbic acid stability during refrigerated storage of orange-carrot juice treated by high pulsed electric field and comparison with pasteurized juice. *J. Food Eng.* **73**:339–345.
- Tuladhar, E., Bouwknecht, M., Zwietering, M. H., Koopmans, M. and Duijzer, E. (2012). Thermal stability of structurally different viruses with proven or potential relevance to food safety. *J. Appl. Microbiol.* **112**:1050–1057.
- Ünal, M. and Bellur, E. (2009). Extraction and characterisation of pectin methylesterase from black carrot (*Daucus carota* L.). *Food Chem.* **116**:836–840.
- U.S. Department of Agriculture, Agricultural Research Service. (1969). Egg pasteurization manual. ARS 74–78. U.S. Department of Agriculture, Agricultural Research Service, Albany, Calif.
- Van Boekel, M. A. J. S. (2008). Kinetic modeling of food quality: A critical review. *Compr. Rev. Food Sci. F.* **7**:144–158.
- Van Buggenhout, S., Sila, D. N., Duvetter, T., Van Loey, A. and Hendrickx, M. (2009). Pectins in Processed Fruits and Vegetables: Part III—Texture Engineering. *Compr. Rev. Food Sci. F.* **8**:105–117.
- Van Duyn, M. A. S. (1999). *Year 2000 Dietary Guidelines: The Case for Fruit and Vegetables First—A Scientific Overview for the Health Professional*. Produce for Better health Foundation, Wilmington, DE.

- Vervoort, L., Van der Plancken, I., Grauwet, T., Verlinde, T., Matser, A., Hendrickx, M. and Van Loey, A. (2012). Thermal versus high pressure processing of carrots: a comparative pilot-scale study on equivalent basis. *Innov. Food Sci. Emerg. Technol.* **15**:1–13.
- Wilbey, R. A. (2014). Heat treatment of foods: principles of pasteurization. *Encyclopedia of Food Microbiology*, 169–174. 2nd edition. Academic Press, San Diego.
- Zang, J., Wang, D. and Zhao, G. (2013). Mechanism of discoloration in processed garlic and onion. *Trends Food Sci. Technol.* **30**:162–173.
- Zhao, L., Wang, S., Liu, F., Dong, P., Huang, W., Xiong, L. and Liao, X. (2013). Comparing the effects of high hydrostatic pressure and thermal pasteurization combined with nisin on the quality of cucumber juice drinks. *Innov. Food Sci. Emerg. Technol.* **17**:27–36.
- Zheng, H. and Lu, H. (2011). Effect of microwave pretreatment on the kinetics of ascorbic acid degradation and peroxidase inactivation in different parts of green asparagus (*Asparagus officinalis* L.) during water blanching. *Food Chem.* **128**:1087–1093.
- Zhou, C., Liu, W., Zhao, J., Yuan, C., Song, Y., Chen, D., Ni, Y. and Li, Q. (2014). The effect of high hydrostatic pressure on the microbiological quality and physical-chemical characteristics of Pumpkin (*Cucurbita maxima* Duch.) during refrigerated storage. *Innov. Food Sci. Emerg. Technol.* **21**:24–34.